

This is the peer reviewed version of the following article: Stojanovic, D., Eyles, S., Cook, H., Alves, F., Webb, M. & Heinsohn, R. (2018) Photosensitive automated doors to exclude small nocturnal predators from nest boxes. *Animal Conservation*, Vol. 22, Iss. 3, Pp 297 - 301; which has been published in final form at <https://doi.org/10.1111/acv.12471>

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

1 **Photosensitive automated doors to exclude small nocturnal predators from nest boxes.**

2

3 Dejan Stojanovic^{1,4}, Sam Eyles², Henry Cook¹, Fernanda Alves³, Matt Webb¹, Robert
4 Heinsohn¹

5 1. Fenner School of Environment & Society, Australian National University, Acton,
6 Australia 2600

7 2. Swift Automation and Mechanical, MacGregor, Australia 2615

8 3. Research School of Biology, Australian National University, Acton, Australia 2600

9 4. Corresponding Author: dejan.stojanovic@anu.edu.au

10 **Running Headline:** A tool to protect nest boxes from predators

11 **Abstract**

12 Nest boxes are a crucial tool for wildlife conservation. Although boxes are often safer from
13 predators than natural nests, if predator and prey are of similar body size survival in boxes
14 may become unacceptably low. Protecting boxes from small predators may be critical to the
15 aims of a project, but no available tools can be reliably deployed for long periods in the field.
16 We trial automated, light sensitive mechanical doors on nest boxes to protect birds nesting
17 in boxes from a small nocturnal predator. At three sites we deployed arrays of nest boxes,
18 and fitted a subset (treatment group) with automated doors, while others were left
19 unprotected. Box occupancy by the target species, clutch size, and nest fate
20 (successful/failed) were monitored using motion activated cameras and by manual checking.
21 Birds in nest boxes fitted with automated doors had a significantly lower risk of nest failure
22 0.25 (\pm 0.11 se) compared to 0.81 (\pm 0.07 se) in the control group. No nests in the treatment

23 group failed due to predation, whereas all nest failures in the control group were
24 attributable to predation. The treatment group did not differ significantly from controls in
25 clutch size. Automated doors operated for a three month breeding season reliably, with
26 minimal maintenance (but battery charge should be monitored). We provide a useful new
27 tool for protecting nest boxes from nocturnal predators, and automated doors did not have
28 any deleterious reproductive consequences on the nests they protected. The automated
29 doors offer practical conservation solutions for nest box conservation programs that (i) are
30 conducted in remote locations with limited accessibility, (ii) require protection measures to
31 be deployed for long periods, (iii) minimise behavioural/physiological impacts on target
32 species, (iv) require targeted protection against nocturnal predators against which more
33 conventional approaches are ineffective or inappropriate.

34 **Key words**

35 Nest predator; nest success; endangered species; conservation biology; nest box; predator
36 protection

37 **1.1 Introduction**

38 Many species are dependent on tree cavities for nesting or shelter sites, but suitable cavities
39 for wildlife can be rare (Gibbons and Lindenmayer, 2002). In some habitats, cavity nesters
40 are limited by the availability of suitable cavities (Newton, 1994), and deforestation
41 exacerbates these shortages (Lindenmayer *et al.*, 2013, Webb *et al.*, 2018). Many cavity
42 nesting species readily occupy artificial nest boxes deployed for research or conservation
43 purposes (Bolton *et al.*, 2004, Flaquer *et al.*, 2006, Olah *et al.*, 2014). Tree cavities passively
44 exclude large predators, making them safe nesting sites (Martin and Pingjun, 1992).
45 Relieving predation pressure may also be an explicit aim of nest box projects (Smith *et al.*,

46 2011). By tailoring nest box design to exclude large predators, survival can be better in nest
47 boxes than natural nests (Bailey and Bonter, 2017, Libois *et al.*, 2012). However, small
48 predators may be able to overcome the passive defence of a small nest cavity entrance hole
49 (Miller, 2002, Stojanovic *et al.*, 2017). Small nocturnal predators of bird nests are globally
50 widespread, and can have important consequences for breeding success (Bradley *et al.*,
51 2003, Williams *et al.*, 2002). In such cases, predation risk in boxes may equal or exceed
52 predation in natural cavities (Evans *et al.*, 2002). In small populations that depend on nest
53 boxes (Stojanovic *et al.*, 2018, Tatayah *et al.*, 2007), small predators pose unacceptable risks
54 to conservation. However the logistic challenges of protecting nests in field settings over a
55 long breeding season remains a major impediment to conservation and ecology projects.

56 In this context, we report the results of a field trial of a new tool for protecting nest boxes.
57 Sugar gliders *Petaurus breviceps* are introduced to Tasmania (Campbell *et al.*, 2018) where
58 they are a major predator of small, tree cavity nesting birds (Stojanovic *et al.*, 2014). There is
59 urgent conservation need to protect birds in nest boxes from sugar gliders (Heinsohn *et al.*,
60 2015). We address this challenge by trialling an automated, solar powered door attached to
61 nest boxes. We use tree martins *Petrochelidon nigricans* to evaluate the efficacy of our
62 automated doors because they are an abundant occupant of nest boxes in our study
63 system. Tree martins are obligate tree cavity nesters and suffer predation from sugar gliders
64 (Stojanovic *et al.*, 2014). Our study aimed to: (1) trial the efficacy of automated doors at
65 protecting bird nests from sugar gliders, and (2) investigate whether operation of the doors
66 impacted key demographic parameters of birds.

67 **2.1 Materials and Methods**

68 We developed and field-tested photosensitive doors for nest boxes (referred to as 'Possum-
69 keeper-outterers' during fund-raising activities, hereafter PKOs). 60 nest boxes were erected
70 at three locations in south eastern Tasmania in December 2017 – Feb 2018 (20 boxes per
71 site). The three sites (Southport Lagoon: S43°28', E146°56'; Meehan Range: S42°49',
72 E147°24', Tooms Lake: S42°13', E147°47') were characterised by dry forests and selected
73 based on high sugar glider predation risk (Heinsohn *et al.*, 2015) and presence of swift
74 parrots *Lathamus discolor* (which are critically endangered by sugar glider predation,
75 Heinsohn *et al.*, 2015), tree martins and sugar gliders at the time of the study. Other
76 potential nocturnal nest predators (e.g. brush-tailed possums *Trichosurus vulpecula*,
77 Tasmanian boobooks *Ninox leucopsis*) and other diurnal nest predators were all present at
78 the time of the study at all sites. Nest boxes occupied by tree martins were randomly
79 assigned to either treatment (up to five nest boxes per site) or control groups (all other nest
80 boxes at the site). Nests were monitored with motion activated cameras (Reconyx™)
81 attached within 20 cm of the nest box entrance hole. PKOs and cameras were deployed on
82 nest boxes after tree martin nest construction began but before the first egg was laid.

83 PKOs incorporate a photosensitive trigger mechanism that causes the door to open/close
84 when ambient light exceeds/falls below 20 lumens (effectively first and last light of each
85 day). This light level was chosen based on a trial of PKOs before the experiment was
86 implemented and using data on first/last nest visitation by swift parrots from motion
87 activated cameras (Stojanovic, D. unpublished data). We opted for a light sensor rather than
88 a clock with fixed open/close schedules because at our high latitude field site, day length
89 varies by ~ 4h/day over the course of a breeding season. PKOs were powered in the field
90 deployments by a 12V28A car battery, recharged continuously by a 12V4A solar panel. Trees
91 with dense canopies that shaded the solar panels were assigned a second panel to

92 compensate. Panels and batteries were deployed in the tree below the nest boxes using
93 5cm external wood screws on straight, unobstructed sections of trunk to protect equipment
94 and cables. PKOs were attached to nest boxes using screws, leaving a gap of ~5 mm
95 between the door and the box face (to prevent snagging). Nest boxes were randomly
96 oriented, so PKOs experienced a range of prevailing weather and light conditions depending
97 on the orientation of the nest box, and which side of the tree the nest box was situated on.
98 Components and assembly instructions for PKOs are provided in Supplementary Materials.

99 To test the efficacy of PKOs at protecting bird nests from sugar glider predation (aim one),
100 we recorded nest fate as successful (at least 1 nestling surviving to fledge) or unsuccessful
101 (no surviving nestlings). Nest fate and confirmation of predation by sugar gliders was
102 determined by reviewing images from the cameras and inspecting nests manually to look
103 for egg fragments and carcasses. We fitted four generalised linear models using nest success
104 as the response variable, with binomial error distributions, and four fixed effects: (i) null, (ii)
105 study site, (iii) treatment type, and (iv) study site + treatment type.

106 To investigate whether PKOs impacted key demographic parameters of birds (aim two) we
107 recorded clutch size of each tree martin nest (as an index of nest productivity and was
108 known for all but one nest). We fitted four generalised linear models using clutch size as the
109 response variable, with Poisson error distributions and the same four fixed effects as above.

110 Competing models were compared using $\Delta AIC < 2$, and all analyses were undertaken in R (R
111 Development Core Team, 2017).

112 **3.1 Results**

113 We recorded 47 tree martin nesting attempts, and 17 of these were successful. Of the 30
114 nests that failed four were in the treatment group, and 26 in the control group (Table 1).

115 **Table 1.** Sample size of tree martin nests per site and treatment group, presented as
116 number of failed nests/total number of nests. * two successive nesting attempts occurred in
117 the same nest box.

Site	Control	Treatment	Total
Southport Lagoon	7/8	0/5	13
Meehan Range	12/15	3/6*	21
Tooms Lake	7/8	1/5	13

118

119 Predation by sugar gliders was the sole cause of nest failure in the control group,
120 determined by detection of carcasses or egg fragments in nest boxes, and confirmed by
121 cameras (Fig. 1). At six treatment nests where sugar gliders were detected (Fig. 1), cameras
122 recorded mean 5.3 unsuccessful predation attempts over the nesting period (median: 3,
123 range: 1 to 14), whereas all control nests failed after a single predation attempt
124 (Supplementary Materials, Video). The best model of nest success included only the
125 treatment type. Nests protected by PKOs had a 0.25 (± 0.11 se) probability of failing
126 compared to 0.81 (± 0.07 se) in the control group. Three of the four nests that did not
127 survive in the treatment group failed for unknown reasons (these nests failed during
128 inclement weather, which may have impacted on nestling survival). The fourth was
129 attributable to a PKO failing to open due to battery failure following several days of cloudy
130 weather and shading of the solar panel. A replacement nesting attempt in that nest box was
131 successful after a second solar panel was added to the system. The other PKOs worked

132 correctly (confirmed with camera images) for the duration of the three month study. PKOs
133 required minimal maintenance (intermittent checks of battery voltage) after initial checking
134 and repositioning of solar panels away from shade to ensure battery charge was being
135 maintained. We also observed brush-tailed possums visiting two nest boxes, and PKOs
136 prevented them from reaching into boxes with their forelimbs or snouts. Black currawongs
137 *Strepera fuliginosa* were also detected at 16 nest boxes during the day, but these predation
138 attempts failed because the box entrances were too small.

139 The best model of clutch size was the null model, and we found no effect of study site or
140 treatment group (Table 2).

141 **Table 2.** Mean \pm standard deviation of clutch size of tree martin nests among the three
142 study sites and two treatment groups.

Site	Control	Treatment
Southport Lagoon	2.7 \pm 1.2	3.3 \pm 0.8
Meehan Range	2.9 \pm 1.8	3.6 \pm 0.5
Tooms Lake	3.4 \pm 0.8	3.4 \pm 0.9

143

144 Cameras recorded occasional repeated opening/closing of PKOs during overcast mornings
145 and evenings. This was corrected by addressing voltage drop in the cables by shortening the
146 length of the wiring between the battery and boxes. PKOs cost approximately \$340 USD per
147 unit (including materials and assembly, batteries, solar panels and tree climbing time).

148

149



150



151



152 **Figure 1.** Automated doors successfully excluded sugar gliders from nest boxes containing
153 tree martin nests despite repeated predation attempts (top). Tree martins had a higher
154 probability of nest success in boxes equipped with PKOs. Sugar gliders could enter nest
155 boxes not fitted with automated doors (bottom).

156 **4.1 Discussion**

157 Protecting animals in nest boxes against predators is a fundamental element of projects that
158 require high survival of the target species. Until now, effective tools to protect nest boxes

159 from small predators have been unavailable despite urgent need. Sugar gliders were
160 unsuccessful despite repeated attempts to prey on nests fitted with PKOs, which improved
161 nest success by 56 % relative to the control group. Predation accounted for all nest failures
162 in boxes without PKOs, and predation events involved the death of adult tree martins and
163 their eggs/nestlings. Our results demonstrate the efficacy of the PKO at eliminating
164 predation even where background predation risk was high and predators persistent. Our
165 results are also encouraging for species vulnerable to larger bodied predators (Beggs and
166 Wilson, 1991), because PKOs prevented brush-tailed possums from reaching into nest
167 boxes, and the design we use could be scaled to suit predators of different sizes. Based on
168 these results, PKOs may be a useful new conservation tool for targeted nest protection
169 against both small and large nocturnal mammals.

170 Clutch size did not differ between the treatment and control groups. Observations of PKOs
171 in operation did not suggest tree martins were distressed by the movement of the door,
172 which was relatively quiet during operation. We did not explicitly test for behavioural
173 change by nest building tree martins after PKOs were deployed on their nests, and this may
174 warrant investigation for species more sensitive to disturbance. We observed no obvious
175 behaviours indicative of distress, and tree martins typically resumed bringing nesting
176 material to boxes within 15 minutes of PKO deployment. Species that may be more sensitive
177 to disturbance could be managed either by (i) pre-emptively deploying PKOs on all nest
178 boxes, or (ii) deploying 'dummy' PKOs on all nest boxes available, before switching to an
179 operational unit when the target species occupies a nest box. This may overcome potential
180 phobia of newly fitted PKOs, leaving animals to tolerate only the opening/closing of the
181 door at first and last light. Replication of this experiment in a predator free habitat may be
182 necessary to detect subtle behavioural/physiological impacts of PKOs, which may have gone

183 undetected in this study because of the high predation rates we recorded. For swift parrots,
184 which are critically endangered by sugar glider predation (Heinsohn *et al.*, 2015), potential
185 behavioural/physiological impacts of PKO function should be identified and weighed against
186 the risk of severe predation mortality (Stojanovic *et al.*, 2014).

187 PKOs represent a new approach for protecting animals in boxes for the duration of (at least)
188 a three month breeding season. Low maintenance tools are key in field programs in remote
189 locations for threatened species and PKOs performed well in this regard. Shading of the
190 solar panels and overcast conditions caused failure of one PKO. Given the unacceptable
191 mortality risk posed by this scenario, we suggest that in shaded habitats or where
192 maintenance checks of PKOs will be infrequent, additional solar panels or backup batteries
193 may be required. Alternatively, where access to field sites is straightforward, regular
194 swapping of batteries may allow solar panels to be dispensed with altogether. However
195 batteries are heavy, and impractical to carry for long distances in the field, which may limit
196 the range of conditions where this approach is viable.

197 PKOs may also be set to open at night and close during the day, to protect nocturnal species
198 from diurnal predators, or to allow nest boxes to be used as a trap for researching nocturnal
199 mammals. Given the effectiveness, simplicity of manufacture, long term reliability, and the
200 ease of deployment on most standard nest box faces, the PKO is a useful new tool that will
201 enable conservation biologists to overcome the substantial risk posed by predators that can
202 breach traditional passive nest box protection measures.

203

204 **Acknowledgements**

205 The work was primarily funded by 1230 supporters of our crowdfunding campaign
206 “Operation PKO”, with additional support from the Australian Government National
207 Environmental Science Program. BirdLife Australia supported the crowdfunding campaign.
208 Thanks to J. Dielenberg, the Eyles family, D. James, K. Presst and T. Watson. The work was
209 undertaken under an Australian National University Animal Ethics Permit 2017/38 and a
210 Tasmanian Government Scientific Permit TFA17144.

211

212 **References**

213 Bailey, R. L. , Bonter, D. N. (2017). Predator guards on nest boxes improve nesting success of birds.

214 *Wildlife Society Bulletin* 41, 434.

215 Beggs, J. R. , Wilson, P. R. (1991). The kaka *Nestor meridionalis*, a new zealand parrot endangered by

216 introduced wasps and mammals. *Biological Conservation* 56, 23.

217 Bolton, M., Medeiros, R., Hothersall, B. , Campos, A. (2004). The use of artificial breeding chambers

218 as a conservation measure for cavity-nesting procellariiform seabirds: A case study of the

219 madeiran storm petrel (*Oceanodroma castro*). *Biological Conservation* 116, 73.

220 Bradley, J. E., Marzluff, J. M. , Thompson Iii, F. R. (2003). Rodents as nest predators: Influences on

221 predatory behavior and consequences to nesting birds. *The Auk* 120, 1180.

222 Campbell, C. D., Sarre, S. D., Stojanovic, D., Gruber, B., Medlock, K., Harris, S., MacDonald, A. J. ,

223 Holleley, C. E. (2018). When is a native species invasive? Incursion of a novel predatory

224 marsupial detected using molecular and historical data. *Diversity and Distributions*, n/a.

225 Evans, M. R., Lank, D. B., Boyd, W. S. , Cooke, F. (2002). A comparison of the characteristics and fate

226 of Barrow's goldeneye and bufflehead nests in nest boxes and natural cavities. *Condor* 104,

227 610.

228 Flaquer, C., Torre, I. , Ruiz-Jarillo, R. (2006). The value of bat-boxes in the conservation of *Pipistrellus*
229 *pygmaeus* in wetland rice paddies. *Biological Conservation* 128, 223.

230 Gibbons, P. , Lindenmayer, D. B. (2002). *Tree hollows and wildlife conservation in Australia*.
231 Melbourne: CSIRO Publishing.

232 Heinsohn, R., Webb, M. H., Lacy, R., Terauds, A., Alderman, R. , Stojanovic, D. (2015). A severe
233 predator-induced decline predicted for endangered, migratory swift parrots (*Lathamus*
234 *discolor*). *Biological Conservation* 186, 75.

235 Libois, E., Gimenez, O., Oro, D., Mínguez, E., Pradel, R. , Sanz-Aguilar, A. (2012). Nest boxes: A
236 successful management tool for the conservation of an endangered seabird. *Biological*
237 *Conservation* 155, 39.

238 Lindenmayer, D. B., Laurance, W. F., Franklin, J. F., Likens, G. E., Banks, S. C., Blanchard, W., Gibbons,
239 P., Ikin, K., Blair, D., McBurney, L., Manning, A. D. , Stein, J. A. R. (2013). New policies for old
240 trees: Averting a global crisis in a keystone ecological structure. *Conservation Letters* 7, 61.

241 Martin, T. E. , Pingjun, L. (1992). Life history traits of open- vs. Cavity-nesting birds. *Ecology* 73, 579.

242 Miller, K. E. (2002). Nesting success of the great crested flycatcher in nest boxes and in tree cavities:
243 Are nest boxes safer from nest predation? *Wilson Bulletin* 114, 179.

244 Newton, I. (1994). The role of nest sites in limiting the numbers of hole-nesting birds: A review.
245 *Biological Conservation* 70, 265.

246 Olah, G., Vigo, G., Heinsohn, R. , Brightsmith, D. J. (2014). Nest site selection and efficacy of artificial
247 nests for breeding success of scarlet macaws *Ara macao macao* in lowland peru. *Journal for*
248 *Nature Conservation* 22, 176.

249 R Development Core Team (2017). R: A language and environment for statistical computing.).
250 Vienna, Austria: R Foundation for Statistical Computing.

251 Smith, R. K., Pullin, A. S., Stewart, G. B. , Sutherland, W. J. (2011). Is nest predator exclusion an
252 effective strategy for enhancing bird populations? *Biological Conservation* 144, 1.

253 Stojanovic, D., Alves, F., Cook, H., Crates, R., Heinsohn, R., Peters, A., Rayner, L., Troy, S. N. , Webb,
254 M. H. (2018). Further knowledge and urgent action required to save orange-bellied parrots
255 from extinction. *Emu - Austral Ornithology* 118, 126.

256 Stojanovic, D., Rayner, L., Webb, M. , Heinsohn, R. (2017). Effect of nest cavity morphology on
257 reproductive success of a critically endangered bird. *Emu - Austral Ornithology*, 1.

258 Stojanovic, D., Webb, M., Alderman, R., Porfirio, L. , Heinsohn, R. (2014). Discovery of a novel
259 predator reveals extreme but highly variable mortality for an endangered bird. *Diversity and*
260 *Distributions* 20, 1200.

261 Tatayah, R. V. V., Malham, J., Haverson, P. , Van de Wetering, J. (2007). Design and provision of nest
262 boxes for echo parakeets *Psittacula eques* in black river gorges national park, Mauritius.
263 *Conservation Evidence* 4, 16.

264 Webb, M. H., Stojanovic, D. , Heinsohn, R. (2018). Policy failure and conservation paralysis for the
265 critically endangered swift parrot. *Pacific Conservation Biology*.

266 Williams, G. E., Wood, P. B. , Thompson Iii, F. (2002). Are traditional methods of determining nest
267 predators and nest fates reliable? An experiment with wood thrushes (*Hylocichla mustelina*)
268 using miniature video cameras. *The Auk* 119, 1126.

269