



ELSEVIER

Contents lists available at ScienceDirect

Journal of Environmental Management

journal homepage: www.elsevier.com/locate/jenvman

Review

How to improve threatened species management: An Australian perspective

B.C. Scheele^{a,b,*}, S. Legge^{a,b}, D.P. Armstrong^c, P. Copley^d, N. Robinson^{a,b}, D. Southwell^{b,e}, M.J. Westgate^a, D.B. Lindenmayer^{a,b}^a Fenner School of Environment and Society, The Australian National University, Canberra, Australia^b National Environmental Science Programme, Threatened Species Recovery Hub, Australia^c Wildlife Ecology Group, Massey University, Palmerston North, New Zealand^d Parks and Regions, Department of Environment, Water and Natural Resources, South Australia, Australia^e Quantitative and Applied Ecology Group, School of BioSciences, University of Melbourne, Victoria, Australia

ARTICLE INFO

Keywords:

Adaptive management

Conservation

Endangered species management

Invasive species

ABSTRACT

Targeted threatened species management is a central component of efforts to prevent species extinction. Despite the development of a range of management frameworks to improve conservation outcomes over the past decade, threatened species management is still commonly characterised as ad hoc. Although there are notable successes, many management programs are ineffective, with relatively few species experiencing improvements in their conservation status. We identify underlying factors that commonly lead to ineffective and inefficient management. Drawing attention to some of the key challenges, and suggesting ways forward, may lead to improved management effectiveness and better conservation outcomes. We highlight six key areas where improvements are needed: 1) stakeholder engagement and communication; 2) fostering strong leadership and the development of achievable long-term goals; 3) knowledge of target species' biology and threats, particularly focusing on filling knowledge gaps that impede management, while noting that in many cases there will be a need for conservation management to proceed initially despite knowledge gaps; 4) setting objectives with measurable outcomes; 5) strategic monitoring to evaluate management effectiveness; and 6) greater accountability for species declines and failure to recover species to ensure timely action and guard against complacency. We demonstrate the importance of these six key areas by providing examples of innovative approaches leading to successful species management. We also discuss overarching factors outside the realm of management influence that can help or impede conservation success. Clear recognition of factors that make species' management more straightforward – or more challenging – is important for setting realistic management objectives, outlining strategic action, and prioritising resources. We also highlight the need to more clearly demonstrate the benefit of current investment, and communicate that the risk of under-investment is species extinctions. Together, improvements in conservation practice, along with increased resource allocation and re-evaluation of the prioritisation of competing interests that threaten species, will help enhance conservation outcomes for threatened species.

1. Introduction

Threatened species management, based on assessments of species extinction risk, threat identification, prioritisation of species for management, and implementation of targeted management actions, is central to curbing biodiversity loss (Primack, 2006). Despite substantial efforts, notable success has been achieved for relatively few species, and as a result, few threatened species have recovered sufficiently following management interventions to allow delisting (Bottrill et al., 2011; Male and Bean, 2005). In part, this failure can be attributed to a severe lack of resources (Evans et al., 2016; McCarthy et al., 2012). For example in Australia, McCarthy et al. (2008) found that funding for threatened

birds was inadequate to prevent further extinctions and facilitate recovery for most listed species, but also that a relatively small increase in resourcing could substantially improve the conservation status of many species. Australian environmental spending is disproportionately low, with Australia one of only several developed countries featuring in the top 40 underfunded countries for conservation spending (Waldron et al., 2013). Further, there has been a sharp reduction in funding over the past decade, with less than five cents for every \$100 of government spending in 2018 directed to biodiversity conservation (ACF, 2018). However, there are several other prominent reasons for the worsening conservation status of many threatened species, such as a failure to address major threats (Johnson et al., 2017), poor enforcement of

* Corresponding author. Building 141, Fenner School of Environment and Society, Linnaeus Way, Acton, ACT 2601, Australia.
E-mail address: ben.scheele@anu.edu.au (B.C. Scheele).

existing legal protections (Harrison et al., 2016), increasing ignorance of scientific evidence (Sutherland and Wordley, 2017), and a culture of apathy (Russell-Smith et al., 2015). In combination, these issues contribute to inaction or inefficient last minute attempts to rescue species on the brink of extinction (Woinarski et al., 2017).

While part of the blame for ongoing species declines can be attributed to funding shortfalls or socio-political issues, in practice, species declines could also be halted by improving the effectiveness of on-ground conservation effort (Sunderland et al., 2009; Toomey et al., 2017). Although there are examples of successful recovery efforts (see Garnett et al., 2018), many other projects are marred by ad hoc and inefficient planning and implementation (Ferraro and Pattanayak, 2006; Pullin et al., 2004; Sutherland et al., 2004). Further, the effectiveness of conservation management is often poorly evaluated, making it difficult to assess how effective each action was, or what species trajectories would have been in the absence of management intervention (Cresswell and Murphy, 2016). A range of decision frameworks and tools have emerged from conservation planning research to address these challenges (see review by Schwartz et al. (2018)), and there have been substantial advances in the practice of expert elicitation (Hemming et al., 2018). While the increasing use of decision frameworks and support tools over the past 20 years has contributed to enhanced outcomes, there is still substantial room for improvement (Cook et al., 2010; Ferraro and Pattanayak, 2006).

Here we address challenges that can impede conservation management of threatened species in Australia. We have taken an Australian perspective for several key reasons. These include the fact that Australia supports a highly diverse and endemic range of species and ecosystems, some of which have been recently lost (e.g. 35% of modern global mammal extinctions have occurred in Australia (Woinarski et al., 2015)), and many more that are threatened (see: Australian Government, 2018). Australia also has a large and diverse array of species recovery programs, with some failures but also prominent successes, upon which to learn from and improve (Garnett et al., 2018). Finally, there is a strong tradition of research excellence in conservation and environmental management in Australia, thus there is scope and capacity within the nation to improve conservation management standards (Harrison, 2006).

We focus on what can be done to improve threatened species management under current constraints. We acknowledge the need for societal changes in human values and their interaction with the environment (Abson et al., 2017), along with a substantial funding increase (Johnson et al., 2017; Waldron et al., 2013). We also address other over-arching factors outside the realm of management control that can impede conservation success. We consider that despite there being a range of program management frameworks readily available, a large implementation gap remains. We provide illustrative case studies of innovative approaches leading to successful species management, noting that 'success' is context-specific and that long-term success will often require continuation of current management trajectories. The views outlined below are the result of a three day workshop of conservation practitioners and researchers with long-term experience in threatened species management, where we worked to collectively identify pitfalls that can lead to ineffective and inefficient management. By drawing attention to some of the key challenges, and providing ways forward, we hope to present a perspective that leads to improved conservation effectiveness and better conservation outcomes.

2. Challenges and solutions in threatened species management

Once a decision to manage a threatened species has been made, the management process consists of three broad stages; 1) conceptualisation and planning, 2) management implementation and evaluation, and 3) program evaluation and revision (Fig. 1). These broad categories reflect a standard management cycle (e.g. Schwartz et al., 2018). Below, we identify common challenges that occur across each of these

stages and provide examples of how they have been overcome. In addition, we discuss challenges in two overarching elements that profoundly impact management; stakeholder engagement and communication, and leadership and personnel. Our evaluation is not exhaustive, but rather reflects personal experience with issues that most commonly arise in the context of threatened species management in Australia. We refer the reader to Schwartz et al. (2018) for guidance on decisions support tools and frameworks for conservation management. We provide a range of species-specific case-studies of successful management programs overcoming many of the challenges discussed in the following section (see Supporting Material Table S1 for additional examples).

2.1. Conceptualisation and planning

A key set of interrelated deficiencies in threatened species management can stem from management that is based on inadequate understanding of the target species' ecology or threats (Fig. 1). These deficiencies result in poor problem definition, and a lack of clear and realistic objectives. Although the need for robust understanding of the target species' ecology and threat impacts is well established (Caughley and Gunn, 1996), in reality it is poorly adhered to, and often undervalued. Attaining sufficient ecological knowledge generally requires detailed field work to ascertain, among other things, the target species' distribution, habitat requirements, life history parameters, population trajectories, and threat impacts (Table S1; example 1). These processes need to be understood across the target species' entire distribution (to make informed choices of where to prioritise management), or the proportion of the species' distribution where management efforts will be implemented. Because threat impacts and threat tolerance are shaped by environmental and biotic processes that vary across environmental space (Scheele et al., 2017a), information obtained for one population is not always transferable to other populations. It is also crucial to understand interactions between multiple threats and the capacity for changes in one element of a system to amplify the impacts of other threats. Many present day management practices remain focused on a single threat despite growing scientific understanding of the importance of considering threat interactions (Scheele et al., 2017a).

While there is a clear need for more ecological research, constrained budgets mean that research must be strategic, with a focus on resolving uncertainty that will improve management decisions. The development of a conceptual model of the target system can help identify knowledge gaps and where to focus research. For example, Bode et al. (2017) used expert opinion to develop an ecosystem model linking malleefowl (*Leipoa ocellata*) persistence with abiotic and biotic processes in mallee ecosystems. The model helped managers identify direct and indirect threats facing the species, the likely response of populations when each threat interaction is managed, and the most uncertain threat interactions that require further research.

While inadequate knowledge of a target species' ecology and threat impacts can hamper decisions about management alternatives, in many cases there is a need for conservation management to proceed – or at least be initiated – despite knowledge gaps. In such cases, management is commonly guided by expert judgement. For example, expert judgement informed a decision to undertake aerial phosphite application in 1997 to control *Phytophthora cinnamomi* in key areas in south west Western Australia, despite uncertainty about phytotoxicity. Urgent management was deemed necessary because the rapid spread of *P. cinnamomi* was driving the local extinction of at least 11 threatened plant species. Importantly, the early, incisive intervention that was based on expert judgement was complimented with monitoring and evaluation of management effectiveness. Monitoring revealed that populations of most species have stabilised following phosphite application (Barrett and Yates, 2015), and that phytotoxic effects are minimal in these vegetation systems (Rathbone and Barrett, 2018).

Challenges arising from ecological uncertainty are often




Fig. 1. Common pitfalls and potential solutions for improving threatened species management.

compounded by diverse stakeholder interests, trade-offs, resource limitations and ambiguity surrounding management efficacy. One constructive approach to these complexities is to use formal structured decision making, particularly in cases where there is substantial ecological uncertainty and a need for rapid action (Gregory et al., 2012) (Table S1; example 2). Structured decision making provides a framework to evaluate different alternatives in a transparent way to identify optimal solutions and clearly articulated objectives and actions (Gregory et al., 2012; Runge, 2011; Schwartz et al., 2018). In the absence of a formal decision-making process, ad hoc approaches may be prone to bias, fail to clearly specify objectives and consider all alternatives, or how planned actions will achieve stated objectives.

Under a structured decision making approach, it is critical during the planning stage to differentiate between fundamental objectives (for example, reducing extinction risk for a target species) and means objectives (such as increasing breeding success for a target species by 20%) (Gregory et al., 2012). Development of fundamental objectives requires engagement of all relevant stakeholders (see examples 3–5 in Table S1 for case studies where a range of stakeholders were engaged in developing fundamental objectives and evaluating management options). It is important that means objectives are linked with performance measures throughout the life of a program to help evaluate efficacy and allow for ongoing refinement (Tear et al., 2005). Canessa et al. (2016) provide a good example of objective setting in the context of ex-situ conservation. They elicited 32 alternative management strategies from experts on the threatened spotted tree frog (*Litoria spenceri*) and found the strategy that maximised the chance of a wild population persisting for 20 years at the lowest cost. However, they also explored how the optimal strategy depended on the weighting given to the two objectives (i.e. maximise persistence vs minimise cost). In this case, the optimal strategy was sensitive to which objective was considered more important, highlighting the need to explicitly define the fundamental objective early in a management program.

In addition to planning for chosen management actions, contingency planning is a crucial, yet often-neglected, element of threatened species management. Good risk management and planning for unexpected outcomes is not always incorporated into conservation programs (Lindenmayer et al., 2018). Contingency planning requires clear recognition that failure is a possibility – something that managers, practitioners and researchers are not always willing, or able to admit.

For example, red fox (*Vulpes vulpes*) control was initially highly successful for facilitating the recovery of an assemblage of small to medium-sized mammals in south-western Australia, but after an initial increase in abundance, many species experienced major declines. The declines were thought to be associated with increases in feral cat (*Felis catus*) predation (Marlow et al., 2015), following meso-predator release of cats as a consequence of fox control (Wayne et al., 2017). This case highlights two key lessons. First, high quality monitoring was crucial to identifying secondary declines, and second, the need for management to plan for the unexpected.

2.2. Management implementation and evaluation

In many cases, scientific research needs to be better integrated into management, particularly when there is substantial ecological uncertainty about threat impacts or management effectiveness (Fig. 1). Importantly, research needs to be targeted toward resolving knowledge gaps that will improve management decisions (e.g. Stojanovic et al., 2018). Adaptive management – a more complex extension of structured decision making – provides a powerful framework to formally integrate research into management (Table S1; examples 6 and 7). Under an adaptive management approach, uncertainty is identified and then alternative management actions are specifically designed to aid learning by resolving uncertainty, while simultaneously progressing management objectives (Runge, 2011; Westgate et al., 2013).

Despite its intuitive appeal, there are few robust examples of active adaptive management of threatened species (Westgate et al., 2013) (an exception is the recently commenced malleefowl program Table S1; example 6). However, less stringent adaptive management is often appropriate and is more commonly applied. An example is a landscape-scale fire management experiment designed to inform and facilitate the recovery of the threatened Gouldian finch (*Erythrura gouldiae*) in Australia's tropical savanna (Legge et al., 2015). First, research established a negative association between frequent and intense fire and finch body condition indices. Second, this information was used to develop and then implement a new fire management plan, to which Gouldian finches responded positively. More broadly, barriers to successful adaptive management are well-documented in the literature, with poor communication and a lack of collaborative relationships between managers and researchers commonly cited as a key barrier that needs to be

overcome to facilitate management-research integration (Burbidge et al., 2011). Ensuring conservation agencies have research capacity with expertise to expedite management-research relations internally, and to communicate and collaborate with researchers in other institutions, will go some way to addressing this issue.

Once management actions have been implemented, it is essential to conduct reliable monitoring of population responses within a management framework that requires follow up actions as the monitoring progresses (Table S1; examples 8–10). However, a recent review of threatened species monitoring in Australia found that across taxonomic groups, between 21 and 46% of threatened vertebrate species received no monitoring at all (Legge et al., 2018). Further, where monitoring does occur, it is often poorly designed, typically having low power to detect changes in population abundance, or to quantify the efficacy of management actions (Legge et al., 2018). Poor quality monitoring reflects under-resourcing, but also monitoring is frequently inhibited by poor data management practices, with data ‘stock-piled’ over time without analysis, and meta-data and data frequently lost – an issue that is amplified with personnel changes (Lindenmayer and Likens, 2018). Data loss then exacerbates further problems, such as the absence of baselines from which to measure changes and evaluate management success. Finally, and perhaps most troubling, are cases where species are monitored robustly, but the monitoring program lacks triggers for management interventions, resulting in species being monitored to extinction, like the Christmas Island pipistrelle (*Pipistrellus murrayi*) (Lindenmayer et al., 2013). Such failures highlight the need to link monitoring and management closely, and initiating management without delay.

2.3. Program evaluation and revision

Thorough, large-scale evaluations of threatened species conservation efforts in Australia are lacking, a point raised repeatedly within State of the Environment reports (Cresswell and Murphy, 2016) (for a species-specific example of program evaluation see Table S1; example 11). Reporting on resource investment commonly focuses on where and how money was spent, with little quantification of whether management benefited the target species or met objectives (Fig. 1). For example, measures like kilometres of fencing built, or changes in feral predator bait take, are often reported. However, to properly evaluate whether management has had the desired effects, we need to be able to measure whether actions like fencing or predator baiting were associated with increased abundance of the target species of conservation concern (i.e. the fundamental objective) (Table S1; examples 12 and 13). Inadequate evaluation is related to poorly formulated objectives that lack specific, measurable outcomes. As a first step, we suggest that assessment of management effectiveness is incorporated into funding programs, and mandated as a condition of funding being granted (requiring improved monitoring). However, there may be little political appetite for changes requiring greater accountability of spending if funding bodies are currently rewarded with easily met targets that can be used to demonstrate so-called conservation success. In addition, the fundamental mismatch between biological and political funding timeframes remains an ongoing challenge.

At a broader level, evaluation efforts are inhibited by lack of accountability for species declines and extinctions, and the absence of independent reviews of conservation programs (for a species-specific example see Table S1; example 14). The current Australian system lacks a direct chain of accountability for species declines or extinctions, with Woinarski et al. (2017) highlighting the need for legislation adjustments to address this shortcoming. Further, under Australian legislation, the implementation of national recovery actions is not mandated, which inhibits efforts to improve accountability and ensure timely actions. In contrast, the U.S. Endangered Species Act, provides clear legislative responsibilities for the U.S. Fish and Wildlife Service to prepare and implement recovery plans, and develop policy and guidelines for

recovery plan preparation (Hoekstra et al., 2002).

Periodic independent review can be useful to ensure transparency in decision making and can help guard against complacency, where management slips into a ‘business as usual’ routine, without evaluation of new evidence or approaches. Given the complexities and uncertainty inherent in many threatened species management programs, there is a need for periodic re-evaluation of current practises and the foundational knowledge base (Bottrill et al., 2011; Walsh et al., 2013). Avoiding complacency also requires awareness of the shifting baseline phenomena, whereby gradual deterioration in a species’ status or habitats occurs too slowly to be registered by individuals or monitoring programs (Bilney, 2014). For example, plant species with long generation times typical of many Australian threatened woody shrubs and trees may show significant time delays to extinction, well beyond standard monitoring periods (see: Krauss et al., 2010).

2.4. Stakeholder engagement and communication

The need to involve a wide range of decision makers and stakeholders in threatened species conservation has long been recognised (Garnett et al., 2018). In the context of threatened species management in Australia, improved engagement and communication is needed on multiple levels; conservation practitioners and managers are often intermediaries between decision makers that control resources, and stakeholders who are involved in implementing management, or who are impacted by management actions (Table S1; example 15). In the first case, there is a need to provide convincing and robust justification to decision makers of the value and merits of investing in threatened species conservation. The justification for threatened species conservation is strongest for programs that engage a wide range of stakeholders to ensure all objectives are considered, provide clearly articulated benefits, and describe measurable objectives that can be used to demonstrate the value of investing resources. In the second case, stakeholders who are impacted by threatened species protection or management need to be integrated into decision-making processes, and should be kept informed of developments throughout the life-cycle of the program. The aim of this dialogue is to develop mutually acceptable actions, although compromise on both sides is often needed (Redpath et al., 2013). These actions will be most effective when conservation practitioners and managers build strong relationships with the community of affected stakeholders, and work to ensure that decision making is transparent (Mishra et al., 2017). Strong relationships are needed to overcome the ‘knowing-doing’ gap, whereby we know what to do, but communicating what is needed, and then doing it, is a barrier (Hulme, 2014).

In addition to regular and effective communication with decision makers and stakeholders, communication with the broader community is required to increase understanding of conservation challenges and current management approaches (Table S1; example 16) (Redpath et al., 2013). Ultimately, efforts to increase resource allocation to conservation programs are dependent to a large extent on increasing the value the community places on biodiversity conservation. It is the job of conservation scientists and managers to communicate good news stories of successful programs (Garnett et al., 2018), but also to highlight the major challenges involved, and that the consequence of inaction is species extinction.

2.5. Leadership and personnel

Delivery of successful species conservation programs requires strong leadership and a well-functioning team (Black et al., 2011) (Table S1; example 17). Conversely, poor leadership and dysfunctional teams have been identified as a reason for failure in some conservation programs (Battisti, 2017). A key barrier to the success of threatened species programs is indecisive and overly risk averse leadership. Indecisiveness can result from high levels of uncertainty, poor individual leadership,

or in response to broader drivers such as competing priorities, or a lack of political will to act (Meek et al., 2015). In the cases of many threatened species, there is often an urgent need for action, without which there is an imminent risk of extinction (Table S1; example 18). For example, failure to enact threat reduction or establishment of ex-situ populations played a role in the extinction of three Australian vertebrate species between 2009 and 2014 (Woinarski et al., 2017). Conservation success is influenced by myriad factors, many of which are outside the control of conservation practitioners; however, ensuring good leadership and personnel is something that can often be directly shaped by conservation practitioners, and hence something we need to get right (Black et al., 2011).

A commonality we have often observed in successful programs is the important role of a ‘species champion’. Species champions come from a diverse range of backgrounds, from someone in government who advocates strongly for a species, to private landowners, and passionate community members. Species champions can often overcome legislative and funding shortfalls to achieve successful outcomes. Key traits include a robust understanding of the target species’ ecology and threats, but also strong planning skills and the capacity to build and organise a team – including engaging with experts where needed. Species champions are also innovators, developing new approaches to successfully manage species (Clark and Kellert, 1988). Most important, however, is their capacity to be optimistic, including clearly articulate a positive long-term vision for the successful recovery of the target species (i.e. ‘stretch goals’).

Negative sentiments are common in threatened species management, and it is hard to overstate the value of positive visions for success (Redford et al., 2016). For example, Gillespie et al. (2018) identify that the presence of a species champion was crucial for the long-term success of conservation programs for several critically endangered frog species from south-eastern Australia (Table S1; example 19). Of course, having a program driven by a species champion can create vulnerability if that person becomes no longer involved in the program. As such, there is a need to foster institutional memory through the long-term retention of key players, ensure adequate succession planning, and encourage and facilitate knowledge transfer between multiple players (Holling and Meffe, 1996). While long-term institutional involvement is often key, programs that develop a vision broader than any one institution, are often highly successful (Redford et al., 2016).

3. What makes threatened species management so difficult?

In the preceding section we outlined common challenges in threatened species management. However, there are also over-arching factors outside the sphere of management influence that can impede conservation success. Clear recognition of factors that make a species’ management more straightforward – or more challenging – is important for setting realistic management objectives and prioritising resources. Below we outline some recurring impediments to threatened species management.

3.1. Threatened species management conflicts with other priorities

Threatened species management can directly conflict with competing social and economic priorities. This is especially the case when threats to species are primarily human driven, such as habitat loss associated with land-use practices. An example is the conservation of Leadbeater’s possum (*Gymnobelideus leadbeateri*); a critically endangered species that occurs in the wet mountain ash (*Eucalyptus regnans*) forests of Victoria. Long-term research has identified key threats to the survival of the species; habitat loss and degradation from forestry activities and wildfire (Lindenmayer et al., 2016). In terms of promoting this species’ conservation, the science is clear; we need to set aside large areas of forested land for conservation, thereby removing the key threat of logging (Lindenmayer et al., 2016). The economic

costs of ending logging (including job loss) would be more than offset by economic gains to agriculture, water provisioning, tourism, and carbon storage (Keith et al., 2017). However, this solution remains politically challenging due to the reaction of the forestry industry to such a policy. A similar conflict is also apparent in the Australian Capital Territory where government policy promotes continued urban expansion into grassland and woodland ecosystems that provide critical habitat for several threatened reptiles and invertebrate species (Howland et al., 2016). In both cases, known threats to highly threatened species are perpetuated due to prioritisation of competing interests ahead of those species. Improving conservation outcomes in such cases is highly challenging and will ultimately require decision makers to accept greater responsibility for the fate of threatened species. We suggest that it is crucial to explicitly recognise that current policy choices prioritise economic or social interests that are key threats to species, and clearly acknowledge and communicate that threatened species will be negatively impacted, and at risk of extinction if the status quo is maintained. Subsequently, this may trigger a rethink of societal values and re-analysis of the economic value of competing interests; allowing society to choose whether priorities should be shifted to promote species conservation (Abson et al., 2017). Such rethinking is particularly pertinent in light of evidence showing strong public support for threatened species conservation (McCune et al., 2017).

3.2. Unknown threats and unknown solutions

In some cases, species can experience major declines resulting from an unknown threat, or if effective management actions to combat a threat are not known. Enigmatic declines are increasingly common in a globalised world where new threats can rapidly emerge. An illustrative example is the enigmatic declines of hundreds of frog species, both in Australia and globally, from the 1970s onwards. In Australia, the disappearances of several rainforest frog species was first documented in 1979 (Scheele et al., 2017b). Further declines were observed across eastern Australia in the 1980s and 1990s, but it was not until 1998 that a previously unknown disease – chytridiomycosis – was identified and demonstrated to be the cause of frog declines (Scheele et al., 2017b). Following identification of the threat, it took a further decade to develop management actions, and these remain only partially effective (Scheele et al., 2014). In total, it took over three decades from threat emergence and documentation of initial species declines to the development of conservation actions to combat chytridiomycosis. In the intervening years, seven species of frog likely went extinct and a further six species declined severely and are at high risk of extinction without targeted ex-situ programs (Scheele et al., 2017b). While there is no single, easy method for identifying novel threats and developing effective responses, long-term investment in ecological research and monitoring will build capacity to better unravel emerging causes of decline and develop effective management responses.

3.3. Highly mobile and cryptic species

Coordinating and implementing conservation actions across large spatial scales covering multiple land tenures and government jurisdictions is challenging. This is compounded by the fact that information on the ecology, threats and effectiveness of management actions is particularly lacking for nomadic and migratory species (Webb et al., 2017). An example of a species with such management challenges is the regent honeyeater (*Anthochaera phrygia*). This semi-nomadic species is critically endangered with an estimated population size of only 350–400 adults, distributed across a 600 000 km² range across parts of South Australia, Victoria, New South Wales, Australian Capital Territory and Queensland. This species moves across vast distances in response to flowering of nectar producing trees. Restoration plantings have been identified as a key management action, but deciding where to establish planting is highly problematic due to the species’ transient occurrence,

and the lag between planting and maturity. Similar challenges are also evident for international migratory species that face severe threats in parts of their range that undermine conservation actions in other regions, and for which managers in Australia have limited capacity to influence (Hansen et al., 2018). In addition, many other traits common to threatened species, such as irruptive population dynamics (arid rodent species), highly cryptic behaviour (pygmy bluetongue, *Tiliqua adelaidensis*) and extreme rarity (plains wanderer, *Pedionomus torquatus*; night parrot, *Pezoporos occidentalis*), can make management challenging (Dickman et al., 2018).

3.4. Lack of critical reflection on management practices

For management of threatened species to succeed, it is critical not only that proper evaluation is undertaken to understand which actions work and which fail, but also to reflect on the process by which decisions are taken and how those processes can be systematically improved over time. This is the essence of adaptive management, which provides a powerful framework to undertake threatened species management (Runge, 2011). It is particularly useful in cases where there is uncertainty surrounding the effectiveness of different management approaches, and where actions are implemented repeatedly, providing opportunities for learning to be integrated and management adjusted accordingly (Runge, 2011). However, the term adaptive management is at risk of becoming a panchreston – a term associated with such a wide range of definitions that it becomes almost meaningless. A lack of clear consensus of what is adaptive management, especially amongst practitioners, is highly problematic (Westgate et al., 2013). The label of ‘adaptive management’ is applied liberally to cover very simple ad hoc trial and error management programs (with no/limited formal monitoring), right through to complex programs that adhere to the core principles of adaptive management, like conceptual model development, thorough monitoring and program evaluation and refinement. In practice, the best point of this continuum to apply to any conservation problem is context-specific; sometimes ad hoc trial and error is exactly the right approach to take. However, since adaptive management conveys the sense of ‘best-practice’, programs so-labelled are sometimes perceived to be immune from scrutiny and without need for improvement. Failure to recognise the need to continually improve management hinders progress on species conservation.

4. Key lessons from threatened species management in Australia

Based on several decades of our collective experience addressing a diverse range of management, policy, and research challenges relating to threatened species, we have observed several lessons broadly applicable to threatened species management:

1. We need to learn from failure. This requires a cultural shift from hiding and downplaying failures, to openly acknowledging failure and most importantly, ensuring that we learn from failures to avoid repeating the same mistakes (Redford and Taber, 2000). To enable failures to be reported and evaluated, programs need to realistically evaluate failure risk in the planning stage.
2. In complex situations, there is a need to slow down and take stock of current knowledge, identify uncertainty and develop a structured approach to move forward. Too often, with a perceived urgency for action, complex problems are glossed over, which can later resurface, sometimes to create insurmountable problems (Lindenmayer and Likens, 2018). The development of a conceptual model can help identify key areas of uncertainty. At the same time, it is crucial to ensure that management does not stall due to uncertainty.
3. More rigorous evaluation of management efficacy is critical, along with greater information of threatened species trajectories. This can be achieved through long-term population abundance or occurrence monitoring, or detailed evaluation of target species’ demography,

with subsequent projections of population abundance. Without improved monitoring of management efficacy and species’ population trajectories, progress on refining and improving management actions will be unacceptably slow.

4. The formation of successful species recovery teams requires a diversity of people with complementary skillsets (Clark and Westrum, 1989), and bringing in people with different skills during each stage of the program. For example, specialist facilitators with experience in structured decision making can help with the problem definition stage of planning, while statisticians might play a role designing monitoring so that it has sufficient power to detect management effectiveness. Threatened species recovery is a multifaceted process, and unbalanced teams may fail to deliver successful conservation outcomes.
5. When recovery plans are deemed necessary, it is crucial that they are funded and implemented, with clear accountability for actions outlined and enforced, and thorough evaluation of management effectiveness. Evaluations of recovery plans and associated data should be compiled in a central depository for public access.

5. Conclusion

Threatened species conservation is a cornerstone of efforts to curb global biodiversity loss. A universal challenge in threatened species management remains limited resources (Johnson et al., 2017). Efforts to increase funding require managers and scientists to clearly demonstrate the benefit of current investment, and better communicate that the risk of under-investment is species extinctions. In parallel, there is opportunity for improvement in management practices. We highlight the importance of: 1) inclusive stakeholder engagement and communication; 2) fostering strong leadership and the development of ambitious long-term goals; 3) knowledge of target species’ ecology and threats, particularly focusing on filling knowledge gaps that impede development of effective management; 4) setting objectives with measurable outcomes; 5) strategic monitoring and action evaluation; and 6) greater accountability for species declines to ensure timely action and guard against complacency. Together, good conservation practice, along with increased resource allocation and re-evaluation of the prioritisation of competing interests that threaten species, will help ensure enhanced conservation outcomes for threatened species.

Acknowledgements

Funding from the Australian Government’s National Environmental Science Programme through the Threatened Species Recovery Hub facilitated the workshop and preparation of this article. David Coates contributed substantially to the development of the arguments presented in this article.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.jenvman.2018.06.084>.

References

- Abson, D.J., Fischer, J., Leventon, J., Newig, J., Schomerus, T., Vilsmaier, U., von Wehrden, H., Abernethy, P., Ives, C.D., Jager, N.W., 2017. Leverage points for sustainability transformation. *Ambio* 46, 30–39.
- ACF, 2018. Australian Conservation Foundation. Background Brief: Environment Spending in Australia. https://www.acf.org.au/background_briefs, Accessed date: 10 May 2018.
- Australian Government, 2018. Australian Government. Species Profile and Threats Database: EPBC Act List of Threatened Fauna. <http://www.environment.gov.au/cgi-bin/sprat/public/publicthreatenedlist.pl>, Accessed date: 10 May 2018.
- Barrett, S., Yates, C.J., 2015. Risks to a mountain summit ecosystem with endemic biota in southwestern Australia. *Austral Ecol.* 40, 423–432.
- Battisti, C., 2017. How to make (in) effective conservation projects: look at the internal

- context!. *Anim. Conserv.* 20, 305–307.
- Bilney, R.J., 2014. Poor historical data drive conservation complacency: the case of mammal decline in south-eastern Australian forests. *Austral Ecol.* 39, 875–886.
- Black, S.A., Groombridge, J.J., Jones, C.G., 2011. Leadership and conservation effectiveness: finding a better way to lead. *Conserv. Lett.* 4, 329–339.
- Bode, M., Baker, C.M., Benshemesh, J., Burnard, T., Rumpff, L., Hauser, C.E., Lahoz-Monfort, J.J., Wintle, B.A., 2017. Revealing beliefs: using ensemble ecosystem modelling to extrapolate expert beliefs to novel ecological scenarios. *Methods Ecol. Evol.* 8, 1012–1021.
- Bottrill, M.C., Walsh, J.C., Watson, J.E., Joseph, L.N., Ortega-Argueta, A., Possingham, H.P., 2011. Does recovery planning improve the status of threatened species? *Biol. Conserv.* 144, 1595–1601.
- Burbidge, A.H., Maron, M., Clarke, M.F., Baker, J., Oliver, D.L., Ford, G., 2011. Linking science and practice in ecological research and management: how can we do it better? *Ecol. Manag. Restor.* 12, 54–60.
- Canessa, S., Converse, S.J., West, M., Clemann, N., Gillespie, G., McFadden, M., Silla, A.J., Parris, K.M., McCarthy, M.A., 2016. Planning for ex situ conservation in the face of uncertainty. *Conserv. Biol.* 30, 599–609.
- Caughley, G., Gunn, A., 1996. *Conservation Biology in Theory and Practice*. Blackwell Science, Cambridge, U.S.
- Clark, T.W., Kellert, S.R., 1988. Toward a policy paradigm of the wildlife sciences. *Renew. Resour. J.* 6, 7–16.
- Clark, T.W., Westrum, R., 1989. High-performance teams in wildlife conservation: a species reintroduction and recovery example. *Environ. Manag.* 13, 663–670.
- Cook, C.N., Hockings, M., Carter, R., 2010. Conservation in the dark? The information used to support management decisions. *Front. Ecol. Environ.* 8, 181–186.
- Cresswell, I., Murphy, H., 2016. *Biodiversity*. In: *Australia State of the Environment 2016*. Australian Government Department of the Environment and Energy, Canberra. <http://dx.doi.org/10.4226/94/58b65ac828812>. <https://soe.environment.gov.au/theme/biodiversity>.
- Dickman, C.R., Greenville, A.C., Wardle, G.M., 2018. Determining trends in irruptive desert species. In: Legge, S., Lindenmayer, D.B., Robinson, N.M., Scheele, B.C., Southwell, D.M., Wintle, B.A. (Eds.), *Monitoring Threatened Species and Ecological Communities*. CSIRO Publishing, Melbourne.
- Evans, D.M., Che-Castaldo, J.P., Crouse, D., Davis, F.W., Epanchin-Niell, R., Flather, C.H., Frohlich, R.K., Goble, D.D., Li, Y.W., Male, T.D., 2016. Species Recovery in the United States: Increasing the Effectiveness of the Endangered Species Act. *Issues in Ecology Report Number 20*. Ecological Society of America 27 pp.
- Ferraro, P.J., Pattanayak, S.K., 2006. Money for nothing? A call for empirical evaluation of biodiversity conservation investments. *PLoS Biol.* 4, e105.
- Garnett, S., Woinarski, J., Lindenmayer, D., Latch, P., 2018. *Recovering Australian Threatened Species: A Book of Hope*. CSIRO Publishing, Melbourne.
- Gillespie, G.R., Hunter, D.A., Hollis, G., Scheele, B.C., West, M., 2018. A tale of threatened frogs: demonstrating the value of long-term monitoring. In: Legge, S., Lindenmayer, D.B., Robinson, N.M., Scheele, B.C., Southwell, D.M., Wintle, B.A. (Eds.), *Monitoring Threatened Species and Ecological Communities*. CSIRO Publishing, Melbourne.
- Gregory, R., Failing, L., Harstone, M., Long, G., McDaniels, T., Ohlson, D., 2012. *Structured Decision Making: A Practical Guide to Environmental Management Choices*. John Wiley & Sons.
- Hansen, B.D., Clemens, R.S., Gallo-Cajiao, E., Jackson, M.V., Kingsford, R.T., Maguire, G.S., Maurer, G., Milton, D.A., Rogers, D.L., Weller, D.R., Weston, M.A., Woehler, E.J., Fuller, R.A., 2018. Shorebird monitoring in Australia: a successful long-term collaboration among citizen scientists, governments and researchers. In: Legge, S., Lindenmayer, D.B., Robinson, N.M., Scheele, B.C., Southwell, D.M., Wintle, B.A. (Eds.), *Monitoring Threatened Species and Ecological Communities*. CSIRO Publishing, Melbourne.
- Harrison, A.L., 2006. Who's who in conservation biology—an authorship analysis. *Conserv. Biol.* 20, 652–657.
- Harrison, J.R., Roberts, D.L., Hernandez-Castro, J., 2016. Assessing the extent and nature of wildlife trade on the dark web. *Conserv. Biol.* 30, 900–904.
- Hemming, V., Burgman, M., Hanea, A., McBride, M., Wintle, B., 2018. Structured expert elicitation using the IDEA protocol. *Methods Ecol. Evol.* 9, 169–180.
- Hoekstra, J.M., Clark, J.A., Fagan, W.F., Boersma, P.D., 2002. A comprehensive review of Endangered Species Act recovery plans. *Ecol. Appl.* 12, 630–640.
- Holling, C.S., Meffe, G.K., 1996. Command and control and the pathology of natural resource management. *Conserv. Biol.* 10, 328–337.
- Howland, B.W., Stojanovic, D., Gordon, L.J., Fletcher, D., Snape, M., Stirmemans, I.A., Lindenmayer, D.B., 2016. Habitat preference of the striped legless lizard: implications of grazing by native herbivores and livestock for conservation of grassland biota. *Austral Ecol.* 41, 455–464.
- Hulme, P.E., 2014. Bridging the knowing–doing gap: know-who, know-what, know-why, know-how and know-when. *J. Appl. Ecol.* 51, 1131–1136.
- Johnson, C.N., Balmford, A., Brook, B.W., Buettel, J.C., Galetti, M., Guangchun, L., Wilmshurst, J.M., 2017. Biodiversity losses and conservation responses in the Anthropocene. *Science* 356, 270–275.
- Keith, H., Vardon, M., Stein, J.A., Stein, J.L., Lindenmayer, D., 2017. Ecosystem accounts define explicit and spatial trade-offs for managing natural resources. *Nat. Ecol. Evol.* 1, 1683–1692.
- Krauss, J., Bommarco, R., Guardiola, M., Heikkinen, R.K., Helm, A., Kuussaari, M., Lindborg, R., Öckinger, E., Pärtel, M., Pino, J., 2010. Habitat fragmentation causes immediate and time-delayed biodiversity loss at different trophic levels. *Ecol. Lett.* 13, 597–605.
- Legge, S., Garnett, S., Maute, K., Heathcote, J., Murphy, S., Woinarski, J.C., Astheimer, L., 2015. A landscape-scale, applied fire management experiment promotes recovery of a population of the threatened Gouldian finch, *Erythrura gouldiae*, in Australia's tropical savannas. *PLoS One* 10, e0137997.
- Legge, S., Lindenmayer, D.B., Robinson, N.M., Scheele, B.C., Southwell, D.M., Wintle, B.A., 2018. *Monitoring Threatened Species and Ecological Communities*. CSIRO Publishing, Melbourne.
- Lindenmayer, D.B., Blair, D., McBurney, J., Banks, S.C., 2016. Ignoring the science in failing to conserve a faunal icon—major political, policy and management problems in preventing the extinction of Leadbeater's possum. *Pac. Conserv. Biol.* 21, 257–265.
- Lindenmayer, D.B., Likens, G.E., 2018. *Effective Ecological Monitoring*, second ed. CSIRO Publishing, Melbourne.
- Lindenmayer, D.B., Piggott, M.P., Wintle, B.A., 2013. Counting the books while the library burns: why conservation monitoring programs need a plan for action. *Front. Ecol. Environ.* 11, 549–555.
- Lindenmayer, D.B., Wood, J., MacGregor, C., Foster, C., Scheele, B., Tulloch, A., Barton, P., Banks, S., Robinson, N., Dexter, N., 2018. Conservation conundrums and the challenges of managing unexplained declines of multiple species. *Biol. Conserv.* 221, 279–292.
- Male, T.D., Bean, M.J., 2005. Measuring progress in US endangered species conservation. *Ecol. Lett.* 8, 986–992.
- Marlow, N.J., Thomas, N.D., Williams, A.A., Macmahon, B., Lawson, J., Hitchen, Y., Angus, J., Berry, O., 2015. Cats (*Felis catus*) are more abundant and are the dominant predator of woylies (*Bettongia penicillata*) after sustained fox (*Vulpes vulpes*) control. *Aust. J. Zool.* 63, 18–27.
- McCarthy, D.P., Donald, P.F., Scharlemann, J.P., Buchanan, G.M., Balmford, A., Green, J.M., Bennun, L.A., Burgess, N.D., Fishpool, L.D., Garnett, S.T., 2012. Financial costs of meeting global biodiversity conservation targets: current spending and unmet needs. *Science* 338, 946–949.
- McCarthy, M.A., Thompson, C.J., Garnett, S.T., 2008. Optimal investment in conservation of species. *J. Appl. Ecol.* 45, 1428–1435.
- McCune, J., Carlsson, A.M., Colla, S., Davy, C., Favaro, B., Ford, A.T., Fraser, K.C., Martins, E.G., 2017. Assessing public commitment to endangered species protection: a Canadian case study. *FACETS* 2, 178–194.
- Meek, M.H., Wells, C., Tomalty, K.M., Ashander, J., Cole, E.M., Gille, D.A., Putman, B.J., Rose, J.P., Savoca, M.S., Yamane, L., 2015. Fear of failure in conservation: the problem and potential solutions to aid conservation of extremely small populations. *Biol. Conserv.* 184, 209–217.
- Mishra, C., Young, J.C., Fiechter, M., Rutherford, B., Redpath, S.M., 2017. Building partnerships with communities for biodiversity conservation: lessons from Asian mountains. *J. Appl. Ecol.* 54, 1583–1591.
- Primack, R.B., 2006. *Essentials of Conservation Biology*. Mass: Sinauer Associates, Sunderland.
- Pullin, A.S., Knight, T.M., Stone, D.A., Charman, K., 2004. Do conservation managers use scientific evidence to support their decision-making? *Biol. Conserv.* 119, 245–252.
- Rathbone, D., Barrett, S., 2018. Long-term phosphite application maintains species assemblages, richness and structure of plant communities invaded by *Phytophthora cinnamomi*. *Austral Ecol.* <https://doi.org/10.1111/aec.12574>.
- Redford, K.H., Aune, K., Plumb, G., 2016. Hope is a bison. *Conserv. Biol.* 30, 689–691.
- Redford, K.H., Taber, A., 2000. Writing the wrongs: developing a safe-fail culture in conservation. *Conserv. Biol.* 14, 1567–1568.
- Redpath, S.M., Young, J., Evelyn, A., Adams, W.M., Sutherland, W.J., Whitehouse, A., Amar, A., Lambert, R.A., Linnell, J.D., Watt, A., 2013. Understanding and managing conservation conflicts. *Trends Ecol. Evol.* 28, 100–109.
- Runge, M.C., 2011. An introduction to adaptive management for threatened and endangered species. *J. Fish Wildlife Manage.* 2, 220–233.
- Russell-Smith, J., Lindenmayer, D., Kubiszewski, I., Green, P., Costanza, R., Campbell, A., 2015. Moving beyond evidence-free environmental policy. *Front. Ecol. Environ.* 13, 441–448.
- Scheele, B.C., Foster, C.N., Banks, S.C., Lindenmayer, D.B., 2017a. Niche contractions in declining species: mechanisms and consequences. *Trends Ecol. Evol.* 32, 346–355.
- Scheele, B.C., Hunter, D.A., Grogan, L., Berger, L., Kolby, J., McFadden, M., Marantelli, G., Skerratt, L.F., Driscoll, D.A., 2014. Interventions for reducing extinction risk in chytridiomycosis-threatened amphibians. *Conserv. Biol.* 28, 1195–1205.
- Scheele, B.C., Skerratt, L.F., Grogan, L.F., Hunter, D.A., Clemann, N., McFadden, M., Newell, D., Hoskin, C.J., Gillespie, G.R., Heard, G.W., 2017b. After the epidemic: ongoing declines, stabilizations and recoveries in amphibians afflicted by chytridiomycosis. *Biol. Conserv.* 206, 37–46.
- Schwartz, M.W., Cook, C.N., Pressey, R.L., Pullin, A.S., Runge, M.C., Salafsky, N., Sutherland, W.J., Williamson, M.A., 2018. Decision support frameworks and tools for conservation. *Conserv. Lett.* 11, 1–12.
- Stojanovic, D., Alves, F., Cook, H., Crates, R., Heinsohn, R., Peters, A., Rayner, L., Troy, S.N., Webb, M.H., 2018. Further knowledge and urgent action required to save orange-bellied parrots from extinction. *Emu-Austral Ornithology* 118, 126–134.
- Sutherland, T., Sutherland-Groves, J., Shanley, P., Campbell, B., 2009. Bridging the gap: how can information access and exchange between conservation biologists and field practitioners be improved for better conservation outcomes? *Biotropica* 41, 549–554.
- Sutherland, W.J., Pullin, A.S., Dolman, P.M., Knight, T.M., 2004. The need for evidence-based conservation. *Trends Ecol. Evol.* 19, 305–308.
- Sutherland, W.J., Wordley, C.F., 2017. Evidence complacency hampers conservation. *Nature Ecology & Evolution* 1, 1215.
- Tear, T.H., Kareiva, P., Angermeier, P.L., Comer, P., Czech, B., Kautz, R., Landon, L., Mehlman, D., Murphy, K., Ruckelshaus, M., 2005. How much is enough? The recurrent problem of setting measurable objectives in conservation. *AIBS (Am. Inst. Biol. Bull. Sci.) Bull.* 55, 835–849.
- Toomey, A.H., Knight, A.T., Barlow, J., 2017. Navigating the space between research and implementation in conservation. *Conserv. Lett.* 10, 619–625.
- Waldron, A., Mooers, A.O., Miller, D.C., Nibbelink, N., Redding, D., Kuhn, T.S., Roberts, J.T., Gittleman, J.L., 2013. Targeting global conservation funding to limit immediate biodiversity declines. *Proc. Natl. Acad. Sci. U.S.A.* 110, 12144–12148.

- Walsh, J.C., Watson, J.E., Bottrill, M.C., Joseph, L.N., Possingham, H.P., 2013. Trends and biases in the listing and recovery planning for threatened species: an Australian case study. *Oryx* 47, 134–143.
- Wayne, A.F., Maxwell, M.A., Ward, C.G., Wayne, J.C., Vellios, C.V., Wilson, L.J., 2017. Recoveries and cascading declines of native mammals associated with control of an introduced predator. *J. Mammal.* 98, 489–501.
- Webb, M.H., Terauds, A., Tulloch, A., Bell, P., Stojanovic, D., Heinsohn, R., 2017. The importance of incorporating functional habitats into conservation planning for highly mobile species in dynamic systems. *Conserv. Biol.* 31, 1018–1028.
- Westgate, M.J., Likens, G.E., Lindenmayer, D.B., 2013. Adaptive management of biological systems: a review. *Biol. Conserv.* 158, 128–139.
- Woinarski, J.C., Burbidge, A.A., Harrison, P.L., 2015. Ongoing unraveling of a continental fauna: decline and extinction of Australian mammals since European settlement. *Proc. Natl. Acad. Sci. U. S. A.* 112, 4531–4540.
- Woinarski, J.C., Garnett, S.T., Legge, S.M., Lindenmayer, D.B., 2017. The contribution of policy, law, management, research, and advocacy failings to the recent extinctions of three Australian vertebrate species. *Conserv. Biol.* 31, 13–23.