

This is the peer reviewed version of the following article: Lindenmayer, D.B., McBurney, L., Blair, D., Wood, J & Banks S.C (2018) From unburnt to salvage logged: quantifying bird responses to different levels of disturbance severity, *Journal of Applied Ecology*: Volume 55, Issue 2, pp. 1626-1636., which has been published in final form at <https://doi.org/10.1111/1365-2664.13137>

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

PROFESSOR DAVID B LINDENMAYER (Orcid ID : 0000-0002-4766-4088)

Article type : Research Article

Handling Editor: Sharif Mukul

From unburnt to salvage logged: quantifying bird responses to different levels of disturbance severity

David B. Lindenmayer^{1,2}, Lachlan McBurney^{1,2}, David Blair^{1,2}, Jeff Wood^{1,2}, and Sam C. Banks^{1,2}

¹Fenner School of Environment and Society

The Australian National University

Canberra, ACT, 2601, Australia

²Long Term Ecological Research Network

Fenner School of Environment and Society

The Australian National University

Canberra, ACT, 2601, Australia

26 **Contact details:** David.Lindenmayer@anu.edu.au

27 **Word Count:** 8049 words

28

29 **Running Head:** Birds and combinations of fire and logging

30

31 **ABSTRACT**

- 32 1. Forests worldwide are increasingly subject to natural and human disturbances,
33 including wildfires and logging of varying intensity and frequency. Understanding
34 how biodiversity responds to different kinds and combinations of natural and human
35 disturbance is critical to enhanced forest management.
- 36 2. We completed an eight-year study of bird responses across a spectrum of disturbance
37 types in Australian Mountain Ash (*Eucalyptus regnans*) forests following wildfires in
38 2009.
- 39 3. We found evidence of a gradient in bird species richness over the study duration. It
40 was highest in unlogged and unburned (least disturbed) sites, decreasing through
41 burnt unlogged forest (subject to high or low intensity fire), lower still in logged
42 forest, and lowest in the most disturbed sites (subject to salvage logging without
43 island retention). Retention of uncut islands within logged areas increased bird species
44 richness above that found in areas that had been clearcut.
- 45 4. The greatest rate of increase per year after disturbance in bird species richness was on
46 sites burnt by high severity fire but which were not subject to any form of logging.
47 The level of disturbance affected the composition of the bird assemblage. Sites that
48 were unlogged and unburned were more likely to support species that were larger,
49 more mobile, and nested at greater heights above the ground.
- 50 5. *Synthesis and applications.* All forms of logging on burned sites impaired recovery in
51 bird species richness relative to sites subject to high severity fire. Alterations in stand
52 structure and plant species composition (and hence modification in bird habitat
53 suitability) due to logging are the most likely reasons for reduced bird species
54 richness and delayed patterns of recovery. This study highlights the importance for
55 native bird species of retaining patches of unlogged forest not only within otherwise
56 clearcut forest, but also in areas that are burned and subject to salvage logging. We

57 therefore suggest that the adoption of retention harvesting be expanded to include
58 stands disturbed by wildfires.

59 **KEYWORDS:** forest birds, Mountain Ash, natural disturbance, salvage logging, south-
60 eastern Australia, variable retention harvest systems, wildfire, fire, species richness

62 INTRODUCTION

63 Disturbances such as fire, windstorms and floods can have a major influence on both
64 natural and human-modified ecosystems and affect the abundance and diversity of species,
65 nutrient and energy cycling, biomass accumulation, hydrological regimes and other key
66 ecosystem processes (Sousa 1984; Swanson *et al.* 2011; Fairman, Nitschke & Bennett 2016;
67 Pulsford, Lindenmayer & Driscoll 2016). The severity, intensity or frequency of natural
68 disturbance regimes can be altered directly and indirectly by human activities (Bradley,
69 Hanson & DellaSala 2016; Parisien *et al.* 2016) such as patterns of land use (Thompson,
70 Spies & Ganio 2007; Cochrane & Laurance 2008; Taylor, McCarthy & Lindenmayer 2014),
71 climate change (Westerling *et al.* 2006; Abatzoglou & Williams 2016) and the establishment
72 of invasive species (Setterfield *et al.* 2010; Johnstone *et al.* 2016; Jones *et al.* 2016). Thus,
73 there can be additive or interactive effects of human and natural disturbances in biodiversity
74 and key ecosystem processes (Buma 2015; Kishchuk *et al.* 2015; Lindenmayer, Thorn &
75 Banks 2017). Understanding how biodiversity responds to different combinations of
76 disturbances is critical to developing prescriptions that underpin the effective management of
77 natural resources (Frelich 2005; Driscoll *et al.* 2010; Leverkus & Castro 2017; Lindenmayer,
78 Thorn & Banks 2017).

79 A potentially severe form of perturbation in forests is salvage logging in which trees
80 damaged by natural disturbance are harvested in an attempt to recover some of their
81 economic value (Cobb *et al.* 2011; Fraver *et al.* 2017; Leverkus & Castro 2017). Salvage
82 logging is widespread and its use is increasing (Thorn *et al.* 2017), likely as a result of the
83 increase in large-scale intensive natural disturbances globally (Seidl *et al.* 2014). There has
84 also been a rapid increase in the number of studies of salvage logging but many lack data on
85 the effects of some important combinations of natural and human disturbance (D'Amato *et al.*
86 2011; Thorn *et al.* 2017; but see Cobb *et al.* 2011; Kishchuk *et al.* 2015). This includes
87 contrasts between salvage logged areas and places subject to conventional harvesting

88 methods such as clearcutting, but particularly lower intensity silvicultural systems like
89 variable retention harvesting (sensu Gustafsson *et al.* 2012; Fedrowitz *et al.* 2014).

90 Here we quantify the response of forest birds across a range of disturbance types,
91 resulting from fire, logging and a combination of both in the wet Mountain Ash (*Eucalyptus*
92 *regnans*) forests of the Central Highlands of Victoria, south-eastern Australia. Large areas of
93 the study region burned in wildfires in 2009. This event, coupled with subsequent post-fire
94 salvage logging operations and ongoing conventional clearcut logging in unburned forest,
95 provided a unique opportunity to establish a comparative study of disturbance effects (Fig. 1;
96 Table 1). Our study design included replicate sites that were: (1) unburned and unlogged, (2)
97 burned at low severity in 2009, (3) burned at high severity in 2009, (4) subject to
98 conventional clearcut logging operations (i.e. unburned stands were clearcut), (5) subject to
99 variable retention harvesting (in which islands of uncut green forest were retained within
100 cutblocks), (6) subject to conventional post-wildfire salvage logging, and, (7) had been
101 salvage logged but with burned islands of forest retained within the cut area (Fig. 1). This last
102 treatment was an extension of the island retention approach typically employed in
103 conventional green forest variable retention harvesting systems (Fedrowitz *et al.* 2014) but
104 applied in a salvage logging context. Our range of treatments therefore facilitated contrasts in
105 bird responses not only between conventional post-fire salvage logging and conventional
106 clearcut logging but also contrasts among sites where variable retention harvesting was
107 deployed in burned versus unburned forest. Our study design also enabled us to determine
108 whether the effects of salvage logging were more substantial than the effects of clearcut
109 logging alone plus the effects of high-severity fire alone. That is, in quantifying the effects of
110 salvage logging on bird biota, we sought evidence for both additive effects of high severity
111 fire and subsequent clearcut logging as well as interactive impacts (sensu Foster *et al.* 2016)
112 between these two kinds of disturbance.

113 We examined four components of bird response – bird species richness, the
114 composition of the bird assemblage, the occurrence of individual bird species, and bird life
115 history attributes (as part of testing performance filtering hypothesis and functional diversity
116 theory; (Mouillot *et al.* 2012; Aubin *et al.* 2016)). We motivated our investigation by posing
117 two key questions to quantify the three components of biotic response:

118 **Question 1.** *Are birds affected by different combinations of natural and human disturbance?*

119 At the outset of this investigation, we postulated that bird species richness and the detection
120 frequency of individual bird species would be lowest in areas subject to conventional salvage

121 logging and highest in unlogged and unburned sites (see Fig. 1). We also postulated that the
122 composition of the bird assemblage would vary between sites subject to different
123 disturbances with some kinds of species (characterized by particular life history traits) being
124 absent from intensively disturbed areas.

125 The broad focus of Question 1 was on cross-sectional contrasts in bird responses
126 between sites subject to different levels of disturbance severity. However, our work entailed
127 documenting changes in bird responses between 2009 and 2016, thereby providing an
128 opportunity to quantify **temporal** patterns on different kinds of sites. Hence, the second
129 question in our investigation was:

130 **Question 2.** *Does the temporal response of bird species richness and individual species vary*
131 *among sites subject to different kinds of disturbance?* We postulated that, relative to unlogged
132 and unburned sites, the recovery of bird species richness would be slowest in salvage logged
133 sites with no island retention that were those most heavily perturbed (Fig. 1). This was
134 because such sites lack critical structural elements (e.g. dead standing trees) and have the
135 most depauperate vegetation communities of all the forest stand categories that we studied
136 (Lindenmayer & Ough 2006; Blair et al. 2016). In our study, we used our unburnt and
137 unlogged sites as ‘benchmarks’ to evaluate the recovery of disturbed sites (Fig. 1). While a
138 small number of bird species in this forest appear to prefer early post-fire conditions (e.g. the
139 Flame Robin *Petroica phoenicea*; (Lindenmayer et al. 2014)), no species occurs exclusively
140 under early successional conditions so our unburnt and unlogged sites were considered an
141 appropriate benchmark for recovery of species richness.

142 MATERIALS AND METHODS

143 *Study area*

144 Our study area was the Mountain Ash forest in the Toolangi, Marysville and
145 Powelltown districts of the Central Highlands of Victoria, south-eastern Australia (Fig. 2).
146 Stand-replacing fire is one of the predominant forms of natural disturbance in Mountain Ash
147 forests leading to stands of broadly uniform age (Smith *et al.* 2016). We constrained our
148 study to one age class of forest – stands that were 70 years old at the time of the 2009
149 wildfires, having regenerated after previous wildfires in 1939. This was to avoid confounding
150 disturbance treatment effects with forest age effects given that different stand ages of
151 Mountain Ash forest support different faunal assemblages, including birds (Loyn 1985;
152 Lindenmayer *et al.* 2009). In addition, 1939 regrowth forest was the most extensive age class

153 in the Mountain Ash ecosystem at the time of the fires (Burns *et al.* 2015) and it is where
154 almost all timber harvesting activity presently takes place (Flint & Fagg 2007). Old growth
155 stands (which are excluded from logging) are now rare in Mountain Ash ecosystems and
156 constitute approximately 1.2% of the extent of this vegetation type in the Central Highlands
157 region of Victoria (Lindenmayer *et al.* 2012).

158 Our study included three levels of fire severity at a site as determined from on-the-
159 ground measurements of vegetation 1-2 months directly following the 2009 fire. Unburned
160 sites were those which were not subject to fire in 2009. Low severity sites were those where
161 the ground was damaged but the understorey and overstorey remained intact. High-severity
162 sites were those in which plants in the ground, shrub and understorey layers were killed and
163 crowns of overstorey trees consumed.

164 *Experimental design and disturbance classes for contrast*

165 The design for this study took advantage of three major studies that ran concurrently
166 from 2009-2016 in which the surveys for birds all employed broadly similar field sampling
167 protocols (by the same field researchers) (see below).

168 The first investigation was a long-term study of the occurrence of birds on sites that
169 were dominated by 1939 regrowth at the time of the 2009 fires. The study included sites
170 burned at high severity in the 2009 fires, sites burnt at low severity in 2009, and sites that
171 remained unburnt in 2009 (Lindenmayer *et al.* 2014) (Table 1).

172 Our second study was a blocked and replicated experiment designed specifically to
173 contrast vertebrate response (including birds) to variable retention harvesting (Lindenmayer
174 *et al.* 2015). The experiment comprised three key treatments in 1939 regrowth forest: **(1)**
175 unlogged forest, **(2)** forest subject to conventional clearcutting (i.e. no island retention), and
176 **(3)** forest subject to variable retention harvesting in which islands of unlogged forest were
177 retained. The treatments were implemented within an experimental block, with the blocking
178 structure replicated, giving a total of 15 sites (Table 1).

179 Our third study was a blocked and replicated salvage logging experiment initiated
180 immediately following the 2009 wildfires. It was designed to have broad parallels with the
181 experiment on variable retention harvesting, except in a post-fire logging setting. The salvage
182 logging experiment comprised 20 sites in three treatments: **(1)** burned but unlogged sites, **(2)**
183 conventionally salvage logged sites (with no stand retention), and **(3)** salvage logged areas
184 with island retention (Table 1).

185 In the case of salvage logged areas, all trees were killed by high-severity fire – in part
186 because salvage logging is restricted (by regulation and codes of practice) only to those areas
187 subject to very high-severity fire. In the case of conventionally clearcut areas, the only green
188 trees remaining within cutover areas are in the retention islands – all standing trees in the
189 remainder of the cutblock were logged (i.e. cut down). The size of conventionally logged and
190 salvage logged cutblocks varied from 15-40 ha (as per Codes of Practices in these forests).
191 The area of Mountain Ash forest burned in the 2009 fires exceeded 72 000 ha (Cruz *et al.*
192 2012).

193 ***Bird survey protocols***

194 We conducted bird surveys annually between 2009 and 2016 with surveys completed
195 in November/early December which is the breeding season for the majority of species and
196 when summer migrants have arrived. Our first surveys in 2009 occurred after disturbance by
197 fire or logging. Our standardised survey protocol was repeated five-minute point interval
198 counts (*sensu* Pyke & Recher 1983; Ralph, Sauer & Droege 1983). For all broad site types in
199 this study and in each year of sampling, we surveyed each site on two different days to
200 account for day effects (Field, Tyre & Possingham 2002). The count on the first day was
201 completed by a different observer than the count on the second day to account for observer
202 heterogeneity (Cunningham *et al.* 1999; Lindenmayer, Wood & MacGregor 2009). We
203 conducted surveys between sunrise and 9.30 am and did not complete counts on days of rain,
204 fog or high winds.

205 For the 54 sites within the long-term monitoring study, we established a 100 metre
206 transect with permanent plots at 0 m, 50 m and 100m points. We did not assume that
207 individual counts at the three points on the same site were independent. We limited our
208 surveys to birds detected within 50 m of a plot point on a given transect. This was to ensure
209 that the birds recorded were within the particular disturbed sampling unit in question.
210 Standardizing the size of the area sampled meant that it was possible to robustly compare
211 counts made across the different studies which together comprised our study (see below).

212 We also established permanent plots in the variable retention harvesting and salvage
213 logging experiments. The variable retention harvesting experiment entailed establishing a
214 permanent plot within a retention island with only birds recorded within 50 m of the centroid
215 of the plot to ensure that only individuals wholly within the island were counted. The islands
216 were a minimum of 50 metres apart (and separated by clearcut forest). There were three plots,

217 one for each of the three islands that had been retained on the cutblock. A similar protocol
218 was used for the salvage logging experiment in which three islands of uncut forest (which
219 had been burned in the preceding wildfire) were retained and a permanent plot was
220 established within each island. Again, only birds within 50 m of the centroid of the plot were
221 recorded to ensure that only individuals wholly within the island were counted. We did not
222 assume that individual counts at the plot points within a site were independent. For sites
223 subject to conventional clearcutting and conventional salvage logging, we positioned our
224 permanent 50 m survey plots in the same spatial configuration as for the logged areas subject
225 to the variable retention harvesting and salvage logging experiments.

226 *Bird life history attributes*

227 We compiled data on bird species traits to explore relationships between species'
228 identities on sites subject to different kinds of disturbances and particular kinds of life-history
229 attributes (see Supporting Information Table S1; Lindenmayer *et al.* 2018). We summarized
230 data on body mass and life history traits (movement, diet, and foraging substrate) (Handbook
231 of Australian and New Zealand Birds 1990-2007; BirdLife Australia 2014). These traits are
232 thought to reflect the ability of species to respond to environmental change (Luck *et al.*
233 2012).

234 *Statistical analyses*

235 We fitted hierarchical generalized linear models (HGLMs) (Lee, Nelder & Pawitan
236 2006) to our data on bird species richness using quasi-Poisson distributions with a
237 logarithmic link and a gamma distribution with a logarithmic link for random site effects (see
238 Bolker *et al.* 2009). We used a logarithmic link for the fixed effects because we considered
239 that the effects would be approximately multiplicative. We used the conjugate distribution for
240 the random effects for ease of computation and interpretation.

241 We used Wald tests to quantify the significance of terms included in the HGLMs. We
242 fitted models which included the disturbance categories as a single factor together with the
243 interaction with the logarithm of the number of years since the 2009 fire plus one, as well as
244 models in which we treated burn severity, the logging treatment and the study identity as
245 separate factors.

246 Our data for individual species were detection frequencies; that is, the number of
247 individual point counts at a site (out of a maximum of six in any given year – 3 plots per site,
248 surveyed twice by a different observer on a different day) in which a given species was

249 recorded. We fitted hierarchical generalized linear models to detection frequency data for
250 individual species. We used a quasi-binomial distribution with a logit link and a beta-
251 distribution with a logit link for the random site effect. This model assumes that fixed effects
252 are additive on the log odds scale and, as for species richness, we used the conjugate
253 distribution for the random effects. We restricted our analyses to the 22 individual species for
254 which there were detections for 40 or more site-year combinations and at least 60 detections
255 in total (Table S2). Species with fewer detections than this had insufficient data to provide
256 reliable results.

257 We used canonical correspondence analysis (ter Braak 1986; Greenacre 2007) to
258 investigate the effects of disturbance and year on the species assemblage. To avoid distortion
259 of our results by relatively rare species, only species with more than 20 detections (N=35)
260 were included in canonical correspondence analysis. We fitted linear regressions of the
261 resulting species scores on a number of bird life history characteristics. We also tested the
262 effect of year and disturbance in the year by disturbance scores using the interaction as the
263 error term.

264 We completed statistical analyses using Genstat for Windows Release 18.2 (VSN
265 International 2015) and R version 3.2.2 (R Core Team 2016).

266 **RESULTS**

267 *Differences in species richness in response to different combinations of disturbance*

268 Our analyses revealed a highly significant ($\chi^2_9 = 206.2, P < 0.001$) gradient in bird
269 species richness with unburned and unlogged sites supporting significantly greater numbers
270 of species ($11.8 \pm 0.45\text{SE}$) relative to sites subject to conventional salvage logging operations
271 ($3.8 \pm 0.43\text{SE}$) (Fig. 3). Values for mean bird species richness in other categories of sites
272 were generally intermediate between the two extremes, with the highest on unlogged sites
273 (*viz*: those subject to low and high severity fire) and lowest where various kinds of logging
274 operations had occurred (Fig. 3). We found no statistical evidence to suggest the impacts of
275 salvage logging on bird species richness were significantly greater than an additive
276 combination of the effects of high severity fire alone plus and clearcut logging alone. That is,
277 our data contained no evidence of a significant interaction between high severity fire and
278 clearcut logging on bird species richness.

279 *Temporal changes in species richness in response to different combinations of disturbance*

280 There were significant ($\chi^2_9 = 87.2, P < 0.001$) differences in the estimated annual rates
281 of change in mean bird species richness for the different disturbances (Fig. 3). There also was
282 evidence of a positive change in mean bird species richness on sites subject to high levels of
283 disturbance (Fig. 3).

284 We found marked interspecific differences in response to the range of disturbances in
285 our study (Fig. 4). The Crescent Honeyeater (*Phylidonyris pyrrhopterus*) (Fig. 4) ($\chi^2_9 = 24.5,$
286 $P = 0.004$), and the Eastern Yellow Robin (*Eopsaltria australis*) (Supporting Information Fig.
287 S1) ($\chi^2_9 = 25.6, P = 0.002$) were among the relatively few individual species which exhibited
288 a pattern of response similar to that identified for mean species richness (i.e. detection
289 frequency was highest on unlogged and unburnt sites and lowest on conventionally salvaged
290 logged sites (see Fig. 2). The detection frequency of the Flame Robin was highest on sites
291 burned at high severity (Fig. 4) ($\chi^2_9 = 55.0, P < 0.001$) whereas it was highest for the Eastern
292 Spinebill (*Acanthorhynchus tenuirostris*) ($\chi^2_9 = 51.0, P < 0.001$) on variable retention logged
293 areas where retained patches remained unburned (Fig. 4). Several individual species were
294 significantly affected by fire, with adverse effects identified for the Brown Thornbill
295 (*Acanthiza pusilla*) ($\chi^2_2 = 10.3, P = 0.006$), Crescent Honeyeater ($\chi^2_2 = 15.4, P < 0.001$),
296 Eastern Spinebill ($\chi^2_2 = 33.3, P < 0.001$), Eastern Whipbird (*Psophodes olivaceus*) ($\chi^2_2 =$
297 $11.9, P = 0.003$), Eastern Yellow Robin ($\chi^2_2 = 14.0, P < 0.001$), Rose Robin (*Petroica rosea*)
298 ($\chi^2_2 = 34.9, P < 0.001$), Rufous Fantail (*Rhipidura rufifrons*) ($\chi^2_2 = 18.8, P < 0.001$) (Fig. S1).
299 More complex effects were found for other species; for example, the detection frequencies of
300 the Silvereye (*Zosterops lateralis*) ($\chi^2_{82} = 15.6, P < 0.001$) and Golden Whistler
301 (*Pachycephala pectoralis*) ($\chi^2_2 = 36.2, P < 0.001$) were highest on severely burned sites and
302 lowest on sites subject to low severity fire (Fig. S1). As in the case of in quantifying salvage
303 logging impacts on bird species richness, we found no evidence of a significant interaction
304 between high severity fire and clearcut logging for any individual bird species.

305 ***Temporal changes in individual species in response to different combinations of*** 306 ***disturbance***

307 We identified marked inter-specific differences in post-disturbance temporal response
308 of bird species (Fig. 4, Fig. S1). The detection frequency of several bird species increased
309 significantly during the eight years of this study including the Brown Thornbill ($\chi^2_1 = 23.1, P$
310 < 0.001), Olive Whistler (*Pachycephala olivacea*) ($\chi^2_1 = 5.8, P = 0.016$), Pilotbird (*Pycnoptilus*
311 *floccosus*) ($\chi^2_1 = 15.4, P < 0.001$), and White-browed Scrub-wren (*Sericornis frontalis*) ($\chi^2_1 =$

312 23.1, $P < 0.001$). The reverse effect was identified for the Rufous Fantail ($\chi_1^2 = 4.2$, P
313 $= 0.041$), and Spotted Pardalote (*Pardalotus punctatus*) (Fig. S1) ($\chi_1^2 = 5.4$, $P = 0.023$) (Fig.
314 S1).

315 ***Response of the bird assemblage to different combinations of disturbance***

316 In the canonical correspondence analysis, the site by year terms accounted for 20% of
317 the variation, and the first two components accounted for 4.5% and 2.0% of the variation
318 respectively. This suggests that factors other than disturbance have a major effect on species
319 composition. A plot of the types of disturbance as represented by the first two components of
320 the canonical correspondence analysis averaged over years was characterised by a gradient
321 from severely burnt sites in the top left to unburnt sites in the bottom right of the diagram
322 (Fig. 5). A second axis from the canonical correspondence analysis represented a somewhat
323 weaker gradient from logged to unlogged forest (Fig. 5). Fig. S2 shows the locations in the
324 first two dimensions for the 35 most common individual bird species and it suggests the
325 composition of the bird assemblage is related primarily to fire. Two species in particular
326 respond positively to fire, the Flame Robin and the Superb Fairy Wren. In addition, we
327 identified significant relationships between the second component of the canonical
328 correspondence analysis and life history attributes which included positive effects on the
329 component scores of nest height ($F_{1,33} = 5.6$, $P = 0.024$), wing length ($F_{1,33} = 5.3$, $P = 0.028$)
330 and dispersal ratio ($F_{1,32} = 5.4$, $P = 0.027$). This analyses indicated that bird species which
331 nested at greater heights above the ground, were larger, or were more mobile were more
332 likely to occur in unlogged forest.

333 **DISCUSSION**

334 ***Bird responses to different combinations of disturbance***

335 The first key question in our study was: *Are birds affected by different combinations*
336 *of natural and human disturbance?* Consistent with predictions at the outset of this
337 investigation (see Fig. 1), we uncovered strong evidence of a gradient in bird species richness
338 congruent with differences in the increasing intensity of disturbance from unlogged, unburnt
339 forest through to conventionally salvage logged forest (Fig. 1 and Fig. 3). Conventionally
340 salvage logged sites supported approximately 25% of the levels of bird species richness
341 found on unlogged, unburned sites with such differences characterizing our study not only at
342 its inception in 2009 but also eight years later (Fig. 3).

343 In comparison with unburned and unlogged sites, we found that levels of bird species
344 richness were highest in areas with increased amounts of the original stand remaining after
345 disturbance, both following fire (i.e. stands burned at low severity support more of the
346 original stand relative to stands burned at high severity), and logging (where variable
347 retention harvesting methods maintain more of the original stand compared to clearcutting)
348 (Fig. 3). Areas subject to clearcutting and salvage logging supported fewer bird species than
349 stands where variable retention harvesting was employed. This result was broadly consistent
350 with the findings of other empirical studies that have explored bird responses across a range
351 of kinds of disturbance (e.g. Barlow *et al.* 2006) as well as global meta-analyses on variable
352 retention harvesting systems which demonstrated that species richness is generally greater
353 with increasingly levels of stand retention (Fedrowitz *et al.* 2014; Thorn *et al.* 2017). Across
354 the spectrum of sites in our study, stands burned at low or high severity supported higher bird
355 species richness than sites subject to variable retention harvesting (including those that were
356 not subsequently burned as well as those that were burned in 2009) (Fig. 3). Thus, our results
357 for bird species richness indicate that all forms of logging reduced bird species richness
358 relative to that quantified for both burned (but unlogged) sites as well as unburned and
359 unlogged sites (Fig. 3).

360 We suggest that the significant reduction in bird species richness on sites subject to
361 the most severe form of perturbation (i.e. conventional salvage logging) is likely due to
362 changes in vegetation plant species composition, loss of diversity and reduction in other key
363 stand structural elements (e.g. large old trees) typically associated with this form of
364 harvesting (e.g. Foster & Orwig 2006; D'Amato *et al.* 2011; Blair *et al.* 2016; Fraver *et al.*
365 2017; Leverkus & Castro 2017).

366 The pattern we identified for species richness was not replicated for the detection
367 frequencies of most individual bird species. Rather, we found evidence of highly species-
368 specific responses to disturbance (Fig. 4, Fig. S1) which highlights the importance of a
369 broader examination of relationships between the intensity of disturbance and bird life history
370 relationships as outlined below.

371 ***Temporal changes in individual species responses to different combinations of disturbance***

372 At the outset of this investigation, we predicted the recovery of bird species richness
373 would be slowest on sites subject to the most severe kinds of disturbance (Fig. 1). We found
374 that the greatest rate of post-disturbance recovery in species richness was on sites subject to

375 high severity fire (Fig. 3). We acknowledge that rates of recovery are not independent of the
376 degree of reduction in species richness after initial disturbance. That is, there will be limited
377 “recovery” on unburnt and unlogged sites because there was never a decline. Rapid recovery
378 in species richness on sites subject to high severity fire is, in part, a function of the substantial
379 initial reduction in species richness at the time of disturbance, although overall richness still
380 did not approach that of unburnt unlogged sites. Part of the explanation for relatively rapid
381 recovery may be related to high-severity fire that leads to dense natural regeneration around
382 dead trees from the previous fire-killed stand (Blair *et al.* 2016; Smith *et al.* 2016). Stands
383 characterized by rapid regeneration in vegetation structure coupled with numerous dead and
384 burned standing trees may, in turn, provide an array of suitable habitat niches for a range of
385 bird species. Notably, such patterns of positive response in bird species richness were not as
386 pronounced on logged sites. This included areas subject to salvage logging (both
387 conventional salvage logging and those subject to salvage logging but with stand retention) as
388 well as sites which had been conventionally clearcut or subject to variable retention
389 harvesting system (Fig. 3). This suggests that all forms of logging impair the rate of bird
390 recovery relative to that quantified for sites subject to high severity fire. Notably, there might
391 there be an upper bound on recovery rate following wildfire if logging in the surrounding
392 burned forest reduces source populations of birds, akin to the landscape trap hypothesis that
393 has been proposed for Mountain Ash forests (Lindenmayer *et al.* 2011). However, detailed
394 medium to long-term source-sink studies would be required to quantify such risks to bird
395 population recovery (if they exist).

396 ***Limitations of the study***

397 We acknowledge that there some limitations to our study. First, limited data prevented
398 us from analyzing results for rare species, although we recognize there are very few bird
399 species of conservation concern in Mountain Ash forests. For example, the Flame Robin
400 (which is an early successional responder in our study system) is under threat in other
401 Australian ecosystems (Montague-Drake, Lindenmayer & Cunningham 2009). A second
402 limitation was that we combined datasets from three different studies. However, we included
403 study identity in the modeling and the same field staff employed similar sampling methods
404 within one forest type and the same aged forest in that forest type.

405 ***Disturbance and bird life history relationships***

406 Consistent with the performance filtering hypothesis and functional diversity theory
407 (Mouillot *et al.* 2012; Aubin *et al.* 2016), we found evidence that disturbance (particularly
408 fire) affected particular functional groups of birds (and therefore the composition of the bird
409 assemblage). Birds which nested at greater heights in the vegetation, or were larger, or were
410 less mobile were more likely to be associated with unburned forest. The reasons for these
411 results remain unclear. However, it is likely that the short regenerating trees are unsuitable for
412 birds that nest at greater heights. In addition, less mobile (e.g. resident) species may take a
413 prolonged period to recolonize intensively perturbed areas from which they have previously
414 been displaced.

415 ***Implications for conservation and management***

416 Our data suggest that both clearcutting and variable retention harvesting have
417 different effects on birds relative to wildfire (including high-severity fire). The most intense
418 forms of disturbance examined (conventional salvage logging with no island retention) led to
419 the most substantial reduction in bird species richness and also impaired post-disturbance
420 recovery in bird species richness (Fig. 3). Earlier work in Mountain Ash forests highlighted
421 the extent to which salvage logging operations can alter potential nesting and foraging habitat
422 for other groups of animals like arboreal marsupials such as through depleting key elements
423 of stand structure like large old hollow-bearing trees (Lindenmayer & Ough 2006) and
424 resprouting understorey plants (e.g. tree ferns) (Blair *et al.* 2016). These impacts suggest a
425 need to limit the amount of salvage logging in the event of future high-severity wildfires in
426 Mountain Ash forests.

427 There have been proposals to increasingly shift away from clearcutting to retention
428 harvesting in many forest types globally (Gustafsson *et al.* 2012). The results of this study
429 suggest that retention harvesting policies and practices need to be extended beyond green
430 (previously undisturbed) forests to include those that are naturally disturbed (e.g. by fire) and
431 potentially subject to salvage logging. In addition, we suggest it is critically important to
432 ensure that burned areas remain unlogged and are included in the design and implementation
433 of reserves so that protected areas capture the variability of forest conditions and bird
434 communities in a manner that allows for recovery from natural disturbances to proceed
435 unimpeded by post-fire timber harvesting (see DellaSala *et al.* 2015; DellaSala *et al.* 2017).

436 **AUTHORS' CONTRIBUTIONS**

437 DBL, LM and DB conceived the ideas, designed methodology, and collected the data; JW
438 and DBL analysed the data; DBL led the writing of the manuscript. All authors contributed
439 critically to the drafts and gave final approval for publication.

440 **ACKNOWLEDGEMENTS**

441 This study was supported by grants from the Australian Research Council, the Government of
442 Victoria (Department of Environment, Land, Water and Planning; Parks Victoria) and the
443 Graeme Wood Foundation. We thank Mason Crane, Dan Florance, Chris MacGregor,
444 Damian Michael, Sachiko Okada and Thea O'Loughlin for assistance in gathering data on
445 birds. Comments from Jos Barlow, Sharif Mukul, Dominick DellaSala and three anonymous
446 referees greatly improved an earlier version of the paper.

447 **DATA ACCESSIBILITY**

448 Data available from the Dryad Digital Repository doi:10.5061/dryad.24t5j04 (Lindenmayer
449 *et al.* 2018).

450 **SUPPORTING INFORMATION**

451 **Table S1.** Summary of the life history and morphological traits used in the analysis, and the
452 relationship of these traits with environmental change.

453 **Table S2.** Number of detections and number of site x year combinations with birds present
454 for the most common species.

455 **Fig. S1.** Estimated effects of disturbance on percentage detection frequency in 2009 and 2016
456 for 18 individual bird species.

457 **Fig. S2.** First two components from canonical correspondence analysis showing locations in
458 multi-dimensional space for the 35 most common bird species.

459 **REFERENCES**

- 460 Abatzoglou, J.T. & Williams, A.P. (2016) Impact of anthropogenic climate change on
461 wildfire across western US forests. *Proceedings of the National Academy of Sciences*,
462 **113**, 11770-11775.
- 463 Aubin, I., Munson, A.D., Cardou, F., Burton, P.J., Isabel, N., Pedlar, J.H. *et al.* (2016) Traits
464 to stay, traits to move: a review of functional traits to assess sensitivity and adaptive
465 capacity of temperate and boreal trees to climate change. *Environmental Review*, **24**,
466 164-186.

- 467 Barlow, J., Peres, C.A., Henriques, L.M., Stouffer, P.C. & Wunderle, J.M. (2006) The
468 responses of understory birds to forest fragmentation, logging and wildfires: An
469 Amazonian synthesis. *Biological Conservation*, **128**, 182-192.
- 470 BirdLife Australia (2014) Birds in Backyards Bird Finder. Available at
471 <http://www.birdsinbackyards.net/finder/all-species>.
- 472 Blair, D., McBurney, L., Banks, S., W., B. & Lindenmayer, D.B. (2016) Disturbance gradient
473 shows logging affects plant functional groups more than fire. *Ecological Applications*,
474 **26**, 2280-2301.
- 475 Bolker, B.M., Brooks, M.E., Clark, C.J., Geange, S.W., Poulsen, J.R., Stevens, M.H.H. *et al.*
476 (2009) Generalized linear mixed models: a practical guide for ecology and evolution.
477 *Trends in Ecology and Evolution*, **245**, 127-113.
- 478 Bradley, C.M., Hanson, C.T. & DellaSala, D.A. (2016) Does increased forest protection
479 correspond to higher fire severity in frequent-fire forests of the western United States?
480 *Ecosphere*, **7**, e01492.
- 481 Buma, B. (2015) Disturbance interactions: characterization, prediction, and the potential for
482 cascading effects. *Ecosphere*, **6**, Art. 70.
- 483 Burns, E.L., Lindenmayer, D.B., Stein, J., Blanchard, W., McBurney, L., Blair, D. *et al.*
484 (2015) Ecosystem assessment of mountain ash forest in the Central Highlands of
485 Victoria, south-eastern Australia. *Austral Ecology*, **40**, 386-399.
- 486 Cobb, T.P., Morissette, J.L., Jacobs, J.M., Koivula, M.J., Spence, J.R. & Langor, D.W.
487 (2011) Effects of postfire salvage logging on deadwood-associated beetles.
488 *Conservation Biology*, **25**, 94-104.
- 489 Cochrane, M.A. & Laurance, W.F. (2008) Synergisms among fire, land use, and climate
490 change in the Amazon. *Ambio*, **37**, 522-527.
- 491 Cruz, M.G., Sullivan, A.L., Gould, J.S., Sims, N.C., Bannister, A.J., Hollis, J.J. *et al.* (2012)
492 Anatomy of a catastrophic wildfire: The Black Saturday Kilmore East fire in Victoria,
493 Australia. *Forest Ecology and Management*, **284**, 269-285.
- 494 Cunningham, R.B., Lindenmayer, D.B., Nix, H.A. & Lindenmayer, B.D. (1999) Quantifying
495 observer heterogeneity in bird counts. *Australian Journal of Ecology*, **24**, 270-277.
- 496 D'Amato, A.W., Fraver, S., Palik, B., Bradford, J. & Patty, L. (2011) Singular and interactive
497 effects of blowdown, salvage logging, and wildfire in sub-boreal pine systems. *Forest
498 Ecology and Management*, **262**, 2070-2078.
- 499 DellaSala, D., Hanson, C., Lindenmayer, D.B. & Furnish, J. (2015) In the aftermath of
500 mixed- and high-severity fire: logging and related actions degrade mixed and high-

501 severity burn areas. *The Ecological Importance of High-Severity Fires: Nature's*
502 *Phoenix*. (eds D. DellaSala & C. Hanson), pp. 313-347. Elsevier, Amsterdam.

503 DellaSala, D.A., Hutto, R.L.H., C.T., Bond, M.L., Ingalsbee, T., Odion, D. & Baker, W.L.
504 (2017) Accommodating mixed-severity fire to restore and maintain ecosystem
505 integrity with a focus on the Sierra Nevada of California, USA. *Fire Ecology*, **13**,
506 148-171.

507 Driscoll, D.A., Lindenmayer, D.B., Bennett, A.F., Bode, M., Bradstock, R.A., Cary, G.J. *et*
508 *al.* (2010) Fire management for biodiversity conservation: Key research questions and
509 our capacity to answer them. *Biological Conservation*, **143**, 1928-1939.

510 Fairman, T.A., Nitschke, C.R. & Bennett, L.T. (2016) Too much, too soon? A review of the
511 effects of increasing wildfire frequency on tree mortality and regeneration in
512 temperate eucalypt forests. *International Journal of Wildland Fire*, **25**, 831-848.

513 Fedrowitz, K.F., Koricheva, J., Baker, S.C., Lindenmayer, D.B., Palik, B., Rosensvald, R. *et*
514 *al.* (2014) Can retention forestry help conserve biodiversity? A meta-analysis. *Journal*
515 *of Applied Ecology*, **51**, 1669-1679.

516 Field, S.A., Tyre, A.J. & Possingham, H.P. (2002) Estimating bird species richness: how
517 should repeat surveys be organized in time? *Austral Ecology*, **27**, 624-629.

518 Flint, A. & Fagg, P. (2007) Mountain Ash in Victoria's State Forests. *Silviculture Reference*
519 *Manual No. 1*. Department of Sustainability and Environment, Melbourne.

520 Foster, C.N., Sato, C.F., Lindenmayer, D.B. & Barton, P.S. (2016) Integrating theory into
521 disturbance interaction experiments to better inform ecosystem management. *Global*
522 *Change Biology*, **22**, 1325-1335.

523 Foster, D.R. & Orwig, D.A. (2006) Preemptive and salvage harvesting of New England
524 forests: when doing nothing is a viable alternative. *Conservation Biology*, **20**, 959-
525 970.

526 Fraver, S., Dodds, K., Kenefic, L., Seymour, R. & Sypitkowski, E. (2017) Forest structure
527 following tornado damage and salvage logging in northern Maine, USA. *Canadian*
528 *Journal of Forest Research*, **47**, 560-564.

529 Frelich, L.E. (2005) *Forest Dynamics and Disturbance Regimes. Studies from Temperate*
530 *Evergreen-Deciduous Forests*. Cambridge University Press, Cambridge, England.

531 Greenacre, M.J. (2007) *Theory and Applications of Correspondence Analysis*, Second Edition
532 edn. Academic Press, Orlando, Florida.

- 533 Gustafsson, L., Baker, S., Bauhus, J., Beese, W., Brodie, A., Kouki, J. *et al.* (2012) Retention
534 forestry to maintain multifunctional forests: a world perspective. *BioScience*, **62**, 633-
535 645.
- 536 Handbook of Australian and New Zealand Birds (1990-2007) *Handbook of Australian, New*
537 *Zealand and Antarctic Birds, volumes 1-7*. Oxford University Press, Melbourne.
- 538 Johnstone, J.F., Allen, C.D., Franklin, J.F., Frelich, L.E., Harvey, B.J., Higuera, P.E. *et al.*
539 (2016) Changing disturbance regimes, ecological memory, and forest resilience.
540 *Frontiers in Ecology and the Environment*, **14**, 369-378.
- 541 Jones, G.M., Gutierrez, R.J., Tempel, D.J., Whitmore, S.A., Berigan, W.J. & Peery, M.Z.
542 (2016) mega-fires: an emerging threate to old-forest species. *Frontiers in Ecology and*
543 *the Environment*, **14**, 300-306.
- 544 Kishchuk, B.E., Thiffault, E., Lorente, M., Quideau, S., Keddy, T. & Sidders, D. (2015)
545 Decadal soil and stand response to fire, harvest, and salvage-logging disturbances in
546 the western boreal mixedwood forest of Alberta, Canada. *Canadian Journal of Forest*
547 *Research*, **45**, 141-152.
- 548 Lee, Y., Nelder, J.A. & Pawitan, Y. (2006) *Generalized Linear Models with Random Effects:*
549 *Unified Analysis via H-Likelihood*. Chapman & Hall/CRC, Boca Raton.
- 550 Leverkus, A. & Castro, J. (2017) An ecosystem services approach to the ecological effects of
551 salvage logging: valuation of seed dispersal. *Ecological Applications*, **27**, 1057-1063.
- 552 Lindenmayer, D.B., Blanchard, W., McBurney, L., Blair, D., Banks, S., Likens, G.E. *et al.*
553 (2012) Interacting factors driving a major loss of large trees with cavities in an iconic
554 forest ecosystem. *PLOS One*, **7**, e41864.
- 555 Lindenmayer, D.B., Blanchard, W., McBurney, L., Blair, D., Banks, S.C., Driscoll, D.A. *et*
556 *al.* (2014) Complex responses of birds to landscape-level fire extent, fire severity and
557 environmental drivers. *Diversity and Distributions*, **20**, 467-477.
- 558 Lindenmayer, D.B., Hobbs, R.J., Likens, G.E., Krebs, C. & Banks, S.C. (2011) Newly
559 discovered landscape traps produce regime shifts in wet forests. *Proceedings of the*
560 *National Academy of Sciences*, **108**, 15887-15891.
- 561 Lindenmayer, D.B., McBurney, L., Blair, D., Wood, J. & Banks, S.C. (2018) Data from:
562 From unburnt to salvage logged: quantifying bird responses to different levels of
563 disturbance severity. *Dryad Digital Repository*, doi:10.5061/dryad.24t5j04
- 564 Lindenmayer, D.B. & Ough, K. (2006) Salvage logging in the montane ash eucalypt forests
565 of the Central Highlands of Victoria and its potential impacts on biodiversity.
566 *Conservation Biology*, **20**, 1005-1015.

- 567 Lindenmayer, D.B., Thorn, S. & Banks, S. (2017) Please do not disturb. A radical change is
568 needed in the way ecosystems are treated after natural disturbance. *Nature Ecology*
569 *and Evolution*, **1**, art 31.
- 570 Lindenmayer, D.B., Wood, J., McBurney, L., Blair, D. & Banks, S.C. (2015) Single large
571 versus several small: The SLOSS debate in the context of bird responses to a variable
572 retention logging experiment. *Forest Ecology and Management*, **339**, 1-10.
- 573 Lindenmayer, D.B., Wood, J. & MacGregor, C. (2009) Do observer differences in bird
574 detection affect inferences from large-scale ecological studies? *Emu*, **109**, 100-106.
- 575 Lindenmayer, D.B., Wood, J., Michael, D., Crane, M., MacGregor, C., Montague-Drake, R.
576 *et al.* (2009) Are gullies best for biodiversity? An empirical examination of Australian
577 wet forest types. *Forest Ecology and Management*, **258**, 169-177.
- 578 Loyn, R.H. (1985) Bird populations in successional forests of Mountain Ash *Eucalyptus*
579 *regnans* in Central Victoria. *Emu*, **85**, 213-230.
- 580 Luck, G., Lavorel, S., McIntyre, S. & Lumb, K. (2012) Improving the application of
581 vertebrate trait-based frameworks to the study of ecosystem services. *Journal of*
582 *Animal Ecology*, **81**, 1065-1076.
- 583 Montague-Drake, R.M., Lindenmayer, D.B. & Cunningham, R.B. (2009) Factors affecting
584 site occupancy by woodland bird species of conservation concern. *Biological*
585 *Conservation*, **142**, 2896-2903.
- 586 Mouillot, D., Graham, N.A., Vileger, S., Mason, N.W. & Bellwood, D.R. (2012) A
587 functional approach reveals community responses to disturbances. *Trends in Ecology*
588 *and Evolution*, **28**, 167-177.
- 589 Parisien, M.-A., Miller, C., Parks, S.A., DeLancey, E.R., Robinne, F.-N. & Flannigan, M.D.
590 (2016) The spatially varying influence of humans on fire probability in North
591 America. *Environmental Research Letters*, **11**, art. 075005.
- 592 Pulsford, S., Lindenmayer, D.B. & Driscoll, D. (2016) A succession of theories: A
593 framework to purge redundancy in post-disturbance theory. *Biological Reviews*, **91**,
594 148-167.
- 595 Pyke, G.H. & Recher, H.F. (1983) Censusing Australian birds: a summary of procedures and
596 a scheme for standardisation of data presentation and storage. *Methods of Censusing*
597 *Birds in Australia* (ed. S.J. Davies), pp. 55-63. Proceedings of a symposium organised
598 by the Zoology section of the ANZAAS and the Western Australian Group of the
599 Royal Australasian Ornithologists Union. Department of Conservation and
600 Environment, Perth.

601 R Core Team (2016) R: A Language and Environment for Statistical Computing. R
602 Foundation for Statistical Computing, Vienna, Austria.

603 Ralph, C.J., Sauer, J.R. & Droege, S. (1983) Monitoring Bird Populations by Point Counts.
604 USDA Forest Service. Pacific Southwest Research Station.

605 Seidl, R., Schelhaas, M.-J., Rammer, W. & Verkerk, P.J. (2014) Increasing forest
606 disturbances in Europe and their impact on carbon storage. *Nature Climate Change*, **4**,
607 806-810.

608 Setterfield, S.A., Rossiter Rachor, N.A., Hutley, L.B., Douglas, M.M. & Williams, R.J.
609 (2010) Turning up the heat: the impacts of *Andropogon gayanus* (gamba grass)
610 invasion on fire behaviour in northern Australian savannas. *Diversity and*
611 *Distributions*, **16**, 854-861.

612 Smith, A.L., Blanchard, W., Blair, D., McBurney, L., Banks, S.C., Driscoll, D.A. *et al.*
613 (2016) The dynamic regeneration niche of a forest following a rare disturbance event.
614 *Diversity and Distributions*, **22**, 457-467.

615 Sousa, W.P. (1984) The role of disturbance in natural communities. *Annual Review of*
616 *Ecology and Systematics*, **15**, 353-391.

617 Swanson, M.E., Franklin, J.F., Beschta, R.L., Crisafulli, C.M., DellaSala, D.A., Hutto, R.L. *et*
618 *al.* (2011) The forgotten stage of forest succession: early-successional ecosystems on
619 forest sites. *Frontiers in Ecology and the Environment*, **9**, 117-125.

620 Taylor, C., McCarthy, M.A. & Lindenmayer, D.B. (2014) Non-linear effects of stand age on
621 fire severity. *Conservation Letters*, **7**, 355-370.

622 ter Braak, C.J.F. (1986) Canonical correspondence analysis: a new eigenvector technique for
623 multivariate direct gradient analysis. *Ecology*, **67**, 1167-1179.

624 Thompson, J.R., Spies, T.A. & Ganio, L.M. (2007) Reburn severity in managed and
625 unmanaged vegetation in a large wildfire. *Proceedings of the National Academy of*
626 *Sciences*, **104**, 10743-10748.

627 Thorn, S., Bassler, C., Burton, P., Cahall, R., Campbell, J.L., Castro, J. *et al.* (2017) Impacts
628 of salvage logging on biodiversity: a meta-analysis. *Journal of Applied Ecology*,
629 doi:10.1111/1365-2664.12945.

630 VSN International (2015) Genstat for Windows 18th Edition. VSN International, Hemel
631 Hempstead, United Kingdom.

632 Westerling, A.L., Hidalgo, H.G., Cayan, D.R. & Swetnam, T.W. (2006) Warming and earlier
633 spring increase western U.S. forest wildfire activity. *Science*, **313**, 940-943.

634

635 **Fig. 1. Part A.** Conceptual model of types of fire and logging-related disturbances and
636 predicted levels of bird species richness in the Mountain Ash forests of the Central Highlands
637 of Victoria, south-eastern Australia. The spectrum of site types include the *de facto*
638 benchmark unlogged and unburned sites (denoted UD) as the least disturbed areas through to
639 our hypothesized most severely disturbed sites (salvage logged [SC]), salvage logged sites
640 with island retention [denoted SI]). Conventional clearcut areas and sites subject to variable
641 retention harvesting are denoted CC and VR, respectively. **Part B.** Postulated temporal
642 responses in bird species richness in relation to different types of fire and logging-related
643 disturbances

644 **Fig. 2.** The location of the study region in the Central Highlands of Victoria, south-eastern
645 Australia.

646 **Fig. 3.** Estimated bird species richness in 2009 and 2016 on sites subject to different kinds of
647 disturbance ranging from sites that were unburned and unlogged (UD) to sites subject to
648 conventional salvage logging following the 2009 wildfire (denoted SC). Other codes for site
649 types are given in Table 1 and Fig. 1. The lines above the bars in the figure show the
650 estimated standard errors of the difference between 2009 and 2016 for each of the
651 disturbances. Bird surveys were completed in each year from 2009 and species richness is
652 shown only for the start (2009) and end (2016) of the study.

653 **Fig. 4.** Estimated effects of disturbance on percentage detection frequency in 2009 and 2016
654 for five individual bird species. Response patterns for other individual species of birds are
655 shown in Fig. S1 in the supplementary material. The lines above the bars in the figure show
656 the estimated standard errors of the difference between 2009 and 2016 for each of the
657 disturbances. Bird surveys were completed in each year from 2009 and a subset of individual
658 species responses are shown only for the start (2009) and end (2016) of the study.

659 **Fig. 5.** First two components from canonical correspondence analysis showing scores (and
660 therefore locations in multi-dimensional space) for the years in this study and ten types of
661 disturbance examined in this study (see Table 1).

662

663


664 **Table 1.** Summary of the broad site types in the study of bird responses to different
 665 combinations of disturbances in the Mountain Ash forests of the Central Highlands of
 666 Victoria, south-eastern Australia. All sites were regrowth from the 1939 wildfires in 2009
 667 when parts of the study region were burned or when they were logged. Standardizing the age
 668 class in the study avoided confounding between stand age effects, biotic responses, and
 669 various kinds and combinations of disturbances. Not all sites could be surveyed in any given
 670 year and the final column in the table provides information on the number of surveys of sites
 671 across the duration of the study.

Site type	Description and associated citations	Abbreviation	No. of sites	No. of site x year combinations
Undisturbed (Unlogged, unburned)	Forest regenerating after the 1939 wildfires which have remained unlogged and unburned since then	UD	26	171
Low severity fire*	1939 regrowth forest that was burned at low severity in the 2009 fire and was not subject to subsequent salvage logging	LS	8	55
High severity fire*	1939 regrowth forest that was burned at high severity in the 2009 fire but was not subject to subsequent salvage logging	HS	20	77
Variable retention harvesting with island retention, no wildfire in 2009	1939 regrowth forest that was subject to variable retention harvesting between 2006 and 2009 in which islands of forest of 0.5-1.5 ha in size were retained with 15-40 ha cutblocks.	VR	4	20


Conventionally clearcut forest, (with no island retention and no wildfire in 2009)	1939 regrowth forest that was subject to clearcutting between 2006 and 2009 and not subject to wildfire in 2009	CC	2	5
Conventionally clearcut forest, high severity fire in 2009	1939 regrowth forest that was subject to clearcutting between 2006 and 2009, but then burned at high severity in 2009 fires	CC+HS	1	4
Variable retention harvesting with island retention, low severity fire in 2009	1939 regrowth forest that was subject to variable retention harvesting between 2006 and 2009 in which islands of forest of 0.5-1 ha in size were retained with 15-40 ha cutblocks but then burned at low severity in the 2009 fires	VR+LS	3	11
Variable retention harvesting with island retention, high severity fire in 2009	1939 regrowth forest that was subject to variable retention harvesting between 2006 and 2009 in which islands of forest of 0.5-1 ha in size were retained with 15-40 ha cutblocks but then burned at high severity in the 2009 fires	VR+HS	5	24

Conventional salvage logging	1939 regrowth forest that was burned at high severity in the 2009 fires and subject to conventional salvage logging in which the entire damaged stand was clearcut	SC	7	40
Modified salvage logging with island retention	1939 regrowth forest that was burned at high severity in 2009 and then subject to modified salvage logging in which islands of burned forest of 0.5-1 ha in size were retained s	SI	13	64


672 *Fire severity was determined from on-the-ground measurements of vegetation 1-2 months
673 directly following the 2009 fire (see text).




Author Manuscript




jpe_13137_f1b.jpg




jpe_13137_f2.jpg



jpe_13137_f3.jpg



jpe_13137_f4.jpg



jpe_13137_f5.jpg