

This is a peer reviewed version of the following article: Tulloch, A., Auerbach, N., Avery-Gomm, S., Bayraktarov, E., Butt, N., Dickman, C.R., Ehmke, G., Fisher, D.O., Grantham, H., Holden, M.H., Lavery, T.H., Leseberg, N.P., Nicholls, M., O'Connor, J., Roberson, L., Smyth, A.K., Stone, Z., Tulloch, V., Turak, E., Wardle, G.M., Watson, J.E.M. (2018) A decision tree for assessing the risks and benefits of publishing biodiversity data, *Nature Ecology & Evolution*, Vol. 2, pp. 1209-1217; which has been published in final form at:

<https://doi.org/10.1038/s41559-018-0608-1>

1

2 A decision tree for assessing the risks and benefits of publishing 3 biodiversity data

4 Ayesha I.T. Tulloch^{1,2,3*}, Nancy Auerbach⁴, Stephanie Avery-Gomm⁴, Elisa Bayraktarov⁴, Nathalie Butt⁴, Chris R.
5 Dickman³, Glenn Ehmke⁵, Diana O. Fisher⁴, Hedley Grantham², Matthew H. Holden⁴, Tyrone H. Lavery⁶,
6 Nicholas P. Leseberg¹, Miles Nicholls⁷, James O'Connor⁵, Leslie Roberson⁴, Anita K. Smyth⁸, Zoe Stone¹,
7 Vivitskaia Tulloch⁴, Eren Turak^{9,10}, Glenda M. Wardle³, James E.M. Watson^{1,2}

8 1. School of Earth and Environmental Sciences, The University of Queensland, St. Lucia Queensland 4072,
9 Australia

10 2. Wildlife Conservation Society, Global Conservation Program, 2300 Southern Boulevard, Bronx NY
11 10460-1068, USA

12 3. Desert Ecology Research Group, School of Life and Environmental Sciences, University of Sydney,
13 Sydney NSW 2006, Australia

14 4. School of Biological Sciences, The University of Queensland, St. Lucia, Brisbane Queensland 4072,
15 Australia

16 5. BirdLife Australia, Melbourne VIC 3053, Australia

17 6. Biodiversity Institute, University of Kansas, Lawrence Kansas 66045, U.S.A.

18 7. Atlas of Living Australia, National Collections & Marine Infrastructure, CSIRO, Acton ACT 2601,
19 Australia

20 8. Terrestrial Ecosystem Research Network (TERN), The University of Adelaide, North Terrace, Adelaide
21 SA 5005, Australia

22 9. NSW Office of Environment and Heritage, Goulburn St, Sydney NSW 2000, Australia

23 10. Australian Museum, 6 College St Sydney NSW 2000, Australia

24

25 ***Author for correspondence:** Ayesha I.T. Tulloch

26 Desert Ecology Research Group, Heydon-Laurence Building A08, School of Life and Environmental Sciences,

27 University of Sydney, Sydney NSW 2006, Australia. ayesha.tulloch@sydney.edu.au

28

29 **Keywords:** data sharing, risk analysis, decision science, data deficient species, conservation biology, research

30 data publication

31 **Inadequate information on the geographical distribution of biodiversity hampers decision making**
32 **for conservation. Major efforts are underway to fill knowledge gaps, but there are increasing**
33 **concerns that publishing the locations of species is dangerous, particularly for species at risk of**
34 **exploitation. While we recognize that well-informed control of location data for highly sensitive**
35 **taxa is necessary to avoid risks, such as poaching or habitat disturbance by recreational visitors, we**
36 **argue that ignoring the benefits of sharing biodiversity data could unnecessarily obstruct**
37 **conservation efforts for species and locations with low risks of exploitation. We provide a decision-**
38 **tree protocol for scientists that systematically considers both the risks of exploitation and potential**
39 **benefits of increased conservation activities. Our protocol helps scientists assess the impacts of**
40 **publishing biodiversity data and aims to enhance conservation opportunities, promote community**
41 **engagement and reduce duplication of survey efforts.**

42

43 Achieving effective conservation relies on accurate knowledge of where species occur to assist with their
44 management¹⁻³. This is particularly true for rare and endangered species that are at risk of extinction. Despite
45 this, one in six IUCN-listed species are considered data deficient, and conservation practitioners routinely face
46 a paucity of primary data on the temporal and spatial distribution of biodiversity^{4,5,6}. Resolving this issue is
47 urgent: without adequate spatially explicit biodiversity data, good management and policy decisions that
48 enable the protection of species and ecosystems may be unachievable⁷⁻⁹.

49 Primary biodiversity data is evidence that associates a species or taxon with a geographic location within a
50 specified time interval. This may include one or more types of evidence: a sighting, a DNA sample, a verified
51 photographic image or traces such as scats, tracks, nests or burrows that that can be attributed to a given
52 taxon with confidence. Primary data may also provide biologically useful information such as age, sex,
53 breeding status, and population abundance. Today there are not just unprecedented online science data
54 services for researchers, conservationists and the public (e.g. wildlife atlases and scientific data repositories
55 such as <http://aekos.org.au>¹⁰), but an increased willingness to share primary biodiversity data (e.g. via citizen

56 science programs such as eBird ¹¹). Further, scientific journals and funding agencies increasingly request
57 transparently archiving research data ¹²⁻¹⁴.

58 Sharing species occurrence information publicly or privately presents a challenge for scientists because it
59 requires balancing potentially difficult and uncertain trade-offs. For example, shortly after their discovery was
60 published, poaching for the pet trade contributed to the local extinction of Chinese Cave Geckos
61 *Goniurosaurus luyi* in Vietnam ¹⁵, prompting calls to not publish primary biodiversity data ¹⁶. In contrast,
62 primary occurrence data shared by researchers in publicly-available databases and within the scientific
63 literature were critical to recent re-assessments of extinction risk for endemic birds in Bolivia and Australia
64 ^{17,18}; which allowed for accurate assessments of extinction status of up to two-thirds of the examined species
65 that otherwise would have been uncertain. To ensure effective conservation informed by the best available
66 knowledge of species distributions and abundances, we must understand the benefits of sharing data and the
67 costs of not sharing data, rather than only the risks as has been the recent focus. Here, we propose a risk
68 management decision protocol that balances potential negative outcomes for species against the conservation
69 benefits of publishing primary occurrence data. By following our decision tree, scientists collecting biodiversity
70 data will be able to ensure that they do not overlook potential conservation opportunities for study species,
71 and that conservation mistakes do not occur through inappropriate release or restriction of data.

72

73 **How are biodiversity data shared?**

74 Data publication is often carefully managed by data authors and custodians to maintain confidentiality and
75 meet jurisdictional laws and national regulations (Supplementary Table 1). Ways of managing the release of
76 data classified as “sensitive” range from publishing precise locations but changing species identifiers to a
77 classification of “restricted” or to a higher taxonomic resolution such as genus or family (if spatial locations are
78 important to share for conservation purposes), to keeping species names accurate but changing locations to
79 mask true spatial coordinates (e.g. by buffering or masking the location), or restricting species location
80 information completely by withholding it from public access (see Supplementary Table 1).

81 The most comprehensive guide on assessing sensitivities around species and required generalisation rules for
82 publishing species locations is provided by the Global Biodiversity Information Facility (GBIF) ¹⁹. GBIF's protocol
83 is to first identify which species are at risk from harm by human activity, and to second assess the impact of
84 this activity on the taxon. These criteria are used to determine whether a species is flagged as sensitive and are
85 then followed by further rules determining the degree of sensitivity. A subsequent rule determines whether
86 release of information will increase the likelihood of harmful impacts on the species. The assessment for
87 whether data should be released considers what level of generalisation or "denaturing" might be required.
88 These range from no restriction for species classed as *low sensitivity*, to increasing restrictions through data
89 generalisation for *low to medium sensitivity* (0.001°), *medium to high sensitivity* (0.01°), and *highly sensitive*
90 *taxa* (0.1°). All location data are withheld if a species is identified as of high biological significance and under
91 high threat ¹⁹. However, no consideration of the benefits of publishing data is made.

92 There are methods of publishing information on where species occur that do not directly release raw species
93 locality data. Many NGO expeditions assess, and publish data, on the biological value of areas to highlight the
94 need for conservation action, e.g., Conservation International's Rapid Assessment Program shares expedition
95 data online to promote awareness of regions with high biodiversity value and high threat ²⁰. Alternatively,
96 species habitat suitability maps can now be published at high resolutions (down to 10m grid-cell size). Such
97 maps, showing locations that have a high probability of containing the species, often include 1) locations
98 where the species occurs and this is known, 2) locations where the species occurs but this is not known, and 3)
99 locations where the species does not occur but can colonise or be translocated if habitat quality is maintained.
100 It is not possible to distinguish between the latter two categories *a priori* so they are typically represented as a
101 combined mapped area (<https://mol.org/species/map/>). As habitat suitability maps are derived from actual
102 species records they are only meaningful and useful if they are produced using precise rather than denatured
103 locations. Hence it is essential that the experts generating these maps have access to full details of the
104 sightings.

105

106 **Benefits of publishing biodiversity data**

107 Here we define data publishing as the release of primary biodiversity data (defined earlier), or products based
108 on these that link a taxon to a location at a given time, to public databases for use by others. In addition to
109 direct conservation benefits, publishing biodiversity data has multiple benefits for researchers and society
110 including research verification, public engagement, stimulation of new/collaborative research, and informing
111 non-researchers about key ecological or conservation issues ²¹⁻²⁵ (Table 1).

112 For species affected primarily by threats such as climate change and habitat loss, if greater availability of
113 biodiversity data enabled more efficient and cost-effective management decisions, the benefits of revealing
114 population locations may outweigh the overall risk of increasing human exploitation of locations²⁶. For
115 instance, habitat loss due to forestry and farming is the most frequent threat to global terrestrial biodiversity
116 ²⁷. Rare species with poorly known distributions are especially likely to have declined from habitat loss, but
117 new populations are often found in unexpected parts of their former ranges ^{28,29}. Any known location data are
118 crucial to protect the remaining habitats of such species through activities such as building accurate species
119 habitat suitability models ³⁰, which can be incorporated into conservation planning and management. Accurate
120 species distribution models built on fine-resolution location data could result in more effective conservation
121 measures because they can lead to investment in conservation at locations where species occur but have not
122 been sighted and locations where species do not occur but can be colonised or translocated. Sharing data is
123 particularly helpful for data deficient species that often slip through the net of regulatory mechanisms due to
124 poor information on where they are and what threatens them ³¹. Ignoring these species in conservation plans
125 risks failing to preserve important locations as well as diversity in ecological traits and evolutionary features of
126 biodiversity ³².

127 Withholding data and records can lead to perverse outcomes for species requiring management to ensure
128 their persistence. For example, where new locations for threatened species remain undiscovered or are
129 destroyed unknowingly in land development, or there is a false impression of range restriction or small
130 population size. If the objective of government conservation agencies, NGO ecologists, scientists and land
131 managers is to minimise the risk of species extinction (Table 1), then sharing data could help indirectly, by
132 improving information on a species' population size or distribution and enabling a more accurate assessment

133 of threat status, or directly, through enabling increased conservation action in known locations. Additionally,
134 agencies that need occurrence data to manage or assess populations may waste limited resources funding
135 redundant data collection.

136

137 **Risks of publishing biodiversity data**

138 Despite recent data sharing initiatives and regulations (Supplementary Table 1), there is evidence that
139 different types of data collectors have varying perceptions of how sharing data could undermine their own
140 objectives (Table 1). Moreover, there is no doubt that poaching of species highly-valued for traditional
141 medicine and recreational hunting has caused species' population declines and even extinctions (e.g., the
142 Javan rhino *Rhinoceros sondaicus*³³⁻³⁵; Table 2). In addition to documented population declines, human access
143 to habitats has caused individual mortality, changes in wildlife behaviour, reduced reproductive rates and
144 habitat disturbance or loss that affect species' ability to persist in their environment^{36,37}. Individual mortality
145 has a greater impact on rare than common species and can cause feedbacks that eventually lead to population
146 declines. Much of the evidence for data publication leading to species declines is anecdotal, with few instances
147 of a direct link between a decline in a population after data on its location being published (Table 2).

148 Many perceived risks of publishing biodiversity data stem from cultural, social or economic objectives rather
149 than conservation objectives (Table 1). For example, many fishers do not share fishing location data because
150 of concerns their data may be used against them to prosecute for violations or lead to fishing restrictions.

151 Many resource managers view their knowledge as private intellectual property and feel that sharing it with
152 others may put them at an economic and social disadvantage²⁶. To achieve a goal of maximising research
153 output³⁸, a research scientist might be concerned about the extra time and cost required to share
154 reproducible data, which could instead be used to publish more papers or write more grants.

155

156 **A balanced decision tree for sharing biodiversity data**


157 *From risks to opportunities to conserve species*

158 A sole focus on the risk to a species fails to consider situations in which the benefits of sharing data outweigh
159 the benefits of not allowing access to biodiversity data. The context of any decision about data publication
160 should not miss opportunities to conserve species, and needs to consider public and private costs such as from
161 redundant surveying effort or the loss of a species. As such, we propose that scientists follow a decision tree
162 that considers the benefits of sharing biodiversity data (Figure 1), which include highlighting species and places
163 of conservation concern (Table 1). In our decision tree, we assess these kinds of benefits against possible risks
164 of sharing data, such as increased pressure on populations (Table 2). Importantly, our protocol considers all
165 relevant threats to the species, and whether conservation mechanisms are either already in place or could be
166 put in place to mitigate or avoid these. A balanced and transparent evaluation of how, not whether, to share
167 biodiversity data requires owners to clarify the risks and likely impacts to a species from data publication, and
168 at the same time help place this information in a decision-making framework that considers actions to reduce
169 risks of harm to species.

170 *Risk management for species at threat of exploitation*

171 Following a risk assessment approach³⁹ to sharing biodiversity data, we agree with other discussions on data
172 sharing¹⁶ that it is first necessary to identify the risk of published locality data enabling (or increasing) access
173 to a species based on how valuable and accessible it is to collectors, poachers, recreational visitors, or other
174 people with interest in the species (Supplementary Figure 1). This will enable those considering publishing
175 spatial biodiversity data to assess the likely harm to the species or population if visitors disturb or exploit it at
176 published localities.

177



178

179 **Figure 1. Decision tree for publishing biodiversity data from monitoring and surveying.** Green, yellow, orange

180 and red boxes indicate data publishing decisions in order of data restrictiveness. Blue-shaded boxes indicate

181 considerations of conservation benefits and actions to mitigate impacts resulting from data sharing. Grey

182 boxes indicate examples from the text and Tables 2 and 3. "IDs" is abbreviation for "identities" (i.e. species'

183 scientific and/or common names). Question marks suggest how to inform particular steps in the tree: ?¹ follow

184 tree according to associated species; ?¹ Consult CITES; ?² Consult IUCN Red List, Recovery plans, National/State

185 threat assessments; ?³ Consult global accessibility maps, local people, government threatened species officers

186 in jurisdiction; ?⁴ Consult conservation evidence and scientific literature; ?⁵ Consult IUCN Red List, conservation

187 evidence and scientific literature, government threatened species officers in jurisdiction.

188

189 Our protocol accounts for various kinds of risk to species from data publication that have been identified in
190 existing ethical data-publication guidelines (Supplementary Table 1). The main risks are increased exploitation
191 for trade or resource use (*ex situ* threats), or disturbance/destruction of habitat due to human access to
192 localities (*in situ* impacts). High *ex situ* value species are those exploited by the wildlife trade or for resources
193 such as food or timber (see CITES Appendix 1 or 2 species; <https://cites.org/eng>). Well-known examples
194 include the African white and black rhinoceros, all elephant species and many fish. Our decision tree also
195 accounts for the fact that risks to some species might be mitigated by conservation measures, such as
196 restricting access to important sites through regulations or physical barriers (e.g. fencing off reserves); actions
197 that might enable public sharing of data. For species where it is not feasible to restrict access, data publication
198 protocols that mask certain characteristics of the data might be used to protect the species identification or
199 location by the public (Supplementary Table 1), although this might restrict the ability of conservation planners
200 and managers to use the data. We suggest in our decision tree that building a high-resolution habitat model
201 with the data would be a sensible way to publish the data whilst ensuring the exact locations of individuals
202 were masked (Figure 1). The full data could be stored securely and granted after request and assessment of
203 motivations, with data protected by a data sharing agreement. For example, a government conservation
204 agency could collate all threatened species occurrence data from researchers licensed to conduct studies on a
205 species, and build a high-resolution map to provide information to the public about the habitat requirements
206 and distribution of species to engage people whilst not providing site-specific occurrence data that would be
207 available under license for researchers. For some species, the risk is so high that both measures (masking
208 locations and restricting access) should be enacted whilst ensuring that monitoring is undertaken to track
209 changes in the species and their threats – this equates to the strictest protocol in our decision tree (Figure 1).
210 One example is fisheries spawning locations, which are almost impossible to restrict access when in
211 international waters, but have high value and a history of over-harvesting (Example 2 in Table 3).
212 For many high-risk species, releasing public data on species occurrence might increase the risk of species
213 decline if locations were previously unknown. If no conservation measures are in place to avoid these declines,
214 our tree suggests that data be desensitised to mask the identities of species but not locations (Figure 1). This
215 would mean the public can still learn that a precise location has conservation value but would not have specific

216 information on which threatened species occur there, reducing the incentive to visit the location. In some
217 cases, however, sharing location data is unlikely to increase the risk of decline, as population information is
218 already in the public domain, or there is poor access to populations. For these populations, we recommend
219 publishing data without restrictions (Figure 1), as additional information could benefit species by improving
220 the ability to track changes in a population or discover new populations in other locations based on improved
221 knowledge about habitat preferences.

222 Some species have higher value *in situ* than *ex situ*, with lower or non-existent market value. Species with high
223 *in situ* value often have high ecotourism value (e.g. whale sharks, rare birds), and may be directly impacted by
224 human disturbance and pathogen exposure associated with human movement into and out of their habitat
225 (e.g. disruption of bird behaviour through electronic bird song playback). Without appropriate conservation
226 measures such as infrastructure or sensitive guidelines for researchers, threatened or rare species with high *in*
227 *situ* value are vulnerable to perturbation by human visitors, and we recommend restricting data in a way that
228 prevents disturbance, for instance through publishing a habitat map instead of raw locations (Figure 1).

229 Many species may not be directly impacted by exploitation for trade or tourism but are still at risk of indirect *in*
230 *situ* impacts if shared data increase visitation to their localities. For example, the surrounding environment
231 could be negatively affected by vegetation disturbance, soil compaction, or introduction of invasive species, or
232 the species might be impacted through associations with other species. These types of disturbances appear
233 minor compared to direct exploitation but can result in a decline in the condition of the surrounding habitat
234 which may harm health and alter behaviour. For example, a fungal species may not be vulnerable to threats of
235 increased disturbance or wildlife trade, but the tree species with which it shares a symbiotic relationship could
236 suffer high exploitation rates resulting from harvesting for its timber. In this case, the vulnerability of the tree
237 population in addition to the fungus should be considered when assessing how to publish data on the
238 occurrence of the fungus (Figure 1).

239 *Maximising data availability to help conservation*

240 Data sharing among researchers, government agencies, NGOs, and citizen science groups will improve our
241 knowledge of population trends and ecology and our ability to protect species from anthropogenic impacts.

242 Many species such as threatened orchids are vulnerable to *in situ* human recreational activities through
243 habitat degradation, irrespective of whether collection activities are restricted^{40,41}. If conservation measures
244 are not in place, increased visitation resulting from the release of new locations for highly-valued recreational
245 species could cause local population declines, and we recommend restricting data to mask either species'
246 identities or localities (Figure 1 and Example 4 of Malaysian *Rafflesia* in Table 3), depending on whether data
247 have value for mitigating threats. In many cases, however, conservation measures have been implemented to
248 avoid or mitigate declines (e.g. the creation of an exclusion zone to eliminate the chance of visitors disturbing
249 the site, see Example 1 of the Australian Night Parrot *Pezoporus occidentalis* in Table 3). In these cases we
250 recommend making occurrence data public to improve conservation, ecological learning and community
251 engagement (Figure 1).

252 When a species' or population's primary threats are neither *in situ* nor *ex situ* direct exploitation or
253 disturbance (see Example 3 of the Vangunu Giant Rat *Uromys vika* in Table 3), we recommend making data
254 public, due to either little known risk of increased visitation to the site, or little chance that visitation would
255 affect population viability (Figure 1). Even when a species has *in situ* value, the risks of increased visitation
256 might be outweighed by the benefits of publishing data. One example is the Critically Endangered West Indian
257 Ocean Coelacanth *Latimeria chalumnae*, an ancient fish thought to have been extinct for 60 million years. In
258 2000, divers observed coelacanths off South Africa's coast, then tagged several individuals⁴². These rare, slow-
259 growing fish are potentially valuable to collectors, but their deep cave habitats are difficult to access and
260 fisheries bycatch poses a much greater threat to survival than poaching⁴³. The location data have been made
261 publicly available, triggering widespread interest among scientists, managers, and the public. This publicity
262 helped create new marine protected areas, fisheries management measures, and a multinational research
263 programme that has generated over US\$6 million in direct government funding, benefitting many additional
264 species in southern Africa (pers comm., A Paterson, South African Institute for Aquatic Biodiversity).

265 In some cases, it may be impossible to decide whether a species has value in the wildlife trade or is vulnerable
266 to visitation disturbance. Until protocols can be updated with new data, we recommend a precautionary
267 approach that restricts data publication, such as that taken in the case of the newly rediscovered, endangered
268 Night Parrot in Australia (Table 3).

269 *Flexibility to adapt to different contexts and changing information*

270 A major challenge of data publication is the evolving nature of restricted data. Lists of “sensitive” species are
271 useful for some data publication protocols (see Supplementary Table 1), but these lists need regular updating
272 to account for changes in conservation status, knowledge and threats and be adopted on a national or global
273 scale. The IUCN Red List is partially revised each year, but local and regional information on threats to species
274 is often poorly mapped and *ad hoc*⁴⁴. Genuine status changes may be rapid and can apply to previously
275 unrestricted species of Least Concern. For example, five of the six most prominent and economically valuable
276 formerly common ash tree species in North America entered the IUCN Red List in 2017 as Critically
277 Endangered, due to huge mortality from an invasive insect, driven by warming climate⁴⁵. Current data
278 restrictions could also be lifted if new conservation actions are implemented, such as the habitat protections
279 for Night Parrots (see above). Decisions to share data should therefore be updated iteratively and quickly.
280 Because the problem of data sharing is complex, our proposed decision tree is not a one-fits-all solution, and
281 we hope that additional inputs by scientists and other stakeholders will enhance its structure and application
282 to diverse decision contexts. Designation of species-specific data sharing rules will need to be adapted to
283 existing pressures found at national or sub-national scales. Users of the decision protocol should also ensure
284 that criteria used to assess existing conservation and policy mechanisms to protect species can detect
285 situations where policy mechanisms or legislation exist but are not implemented. We developed our decision
286 tree based on the objective of maximizing persistence of a species, but the protocol could be adapted to
287 account for additional objectives, such as maximising public engagement in conservation, or conserving whole
288 ecosystems.

289 *Ensuring data re-use and application to conservation decisions*

290 An essential step to promote data sharing and enhance data re-use is to ensure that users know which data
291 exist and are available. Metadata represent the set of instructions or documentation that describe the
292 content, context, quality, structure, and reusability of a data set. In addition to publishing biodiversity data,
293 making public the background metadata is critical, and could be accompanied by a sample of the database to
294 enable potential users to assess if those data are fit for purpose⁴⁶. We present our protocol anticipating that

295 repositories holding biodiversity data will have cybersecurity data administrators managing the security of
296 holdings. Data policies should state repository security so that data submitters can decide whether the
297 repository is trustworthy. As species locality data are found in multiple repositories, we recommend that the
298 appropriate mode of sharing biodiversity data be a species or population attribute rather than an attribute of a
299 given set of data points specified by data authors. This places greater responsibility on researchers to
300 determine how to share data and the decision tree we have proposed should help this.

301 Although acquiring the information needed to walk through our decision tree could sometimes be time
302 consuming and difficult for individual researchers to obtain, all the information needed for applying our
303 decision tree will be available to those evaluating species for the Convention on International Trade in
304 Endangered Species of Wild Fauna and Flora (CITES) or for IUCN Red Listing. Hence it would make sense for the
305 application of our decision tree to be integrated into these evaluation processes as well as national and
306 subnational assessments of species' threat status and updated regularly.

307

308 **Combating illegal species exploitation**

309 Human exploitation of species for trade, resources, or nature-based recreation continues even in locations
310 with few or no scientific studies. Increased use of social media means that the opportunity to manage sensitive
311 information is declining even if we want to restrict it ⁴⁷. The wide range and varied impacts of threats to
312 species mean that researchers and practitioners have an imperative to understand not only where species
313 occur, but also the spread and intensity of both local and off-site threats to species. Despite government
314 agreements such as CITES, illegal resource take (e.g. unreported fishing) and wildlife trade continues, with
315 black market prices ranging from US\$2 for a sea turtle in Mexico ⁴⁸, to US\$31,000 for an Australian black-
316 cockatoo ⁴⁹, or US\$400,000 for a gorilla ⁵⁰. It is important to articulate whether these kinds of threats, driven
317 by *ex situ* markets, are likely to increase when new localities or ecological information on a population are
318 published. In this way, data can be responsibly and appropriately restricted if threats to a species would
319 increase after publishing new localities, or shared without restriction if new data would not affect species'
320 persistence (see Figure 1).

321 Sharing of species information is without doubt critical in building biodiversity knowledge and managing the
322 global extinction crisis. To date, almost all data publication decisions made by governments, societies or
323 individuals focused on the costs of sharing; benefits are never explicitly quantified, making it impossible to
324 extrapolate data restriction decisions to other species, locations or contexts. Our decision protocol for
325 publishing spatial biodiversity data aims at overcoming this inefficiency and enables scientists to better decide
326 how (and when) to publish data responsibly in repositories. The challenge is to share data in a way that avoids
327 perverse outcomes for biodiversity when it is used. In many cases, sharing data will have greater conservation
328 (and educational) benefits than restricting it from use by those wishing to use it to increase community
329 engagement or to promote conservation actions. Above all else, being explicit about what those benefits
330 might be, and weighing them against the likely risks of making data public, will ensure that species are not put
331 in greater danger from new data being released into the public domain.

332

333 **References**

- 334 1 Costello, M. J., May, R. M. & Stork, N. E. Can We Name Earth's Species Before They Go Extinct? *Science*
335 **339**, 413-416, doi:10.1126/science.1230318 (2013).
- 336 2 Boitani, L. *et al.* What spatial data do we need to develop global mammal conservation strategies?
337 *Philosophical Transactions of the Royal Society B: Biological Sciences* **366**, 2623 (2011).
- 338 3 Rondinini, C., Wilson, K. A., Boitani, L., Grantham, H. & Possingham, H. P. Tradeoffs of different types
339 of species occurrence data for use in systematic conservation planning. *Ecol Lett* **9**, 1136-1145,
340 doi:10.1111/j.1461-0248.2006.00970.x (2006).
- 341 4 Whittaker, R. J. *et al.* Conservation Biogeography: assessment and prospect. *Divers Distrib* **11**, 3-23,
342 doi:10.1111/j.1366-9516.2005.00143.x (2005).
- 343 5 Winter, M. *et al.* Patterns and biases in climate change research on amphibians and reptiles: a
344 systematic review. *Roy Soc Open Sci* **3**, doi:10.1098/rsos.160158 (2016).
- 345 6 Feeley, K. J. & Silman, M. R. Keep collecting: accurate species distribution modelling requires more
346 collections than previously thought. *Divers Distrib* **17**, 1132-1140, doi:10.1111/j.1472-
347 4642.2011.00813.x (2011).

348 7 Joppa, L. N. *et al.* Filling in biodiversity threat gaps. *Science* **352**, 416-418, doi:10.1126/science.aaf3565
349 (2016).

350 8 Zeller, D., Froese, R. & Pauly, D. On losing and recovering fisheries and marine science data. *Marine*
351 *Policy* **29**, 69-73, doi:10.1016/j.marpol.2004.02.003 (2005).

352 9 Walsh, J. C., Dicks, L. V. & Sutherland, W. J. The effect of scientific evidence on conservation
353 practitioners' management decisions. *Conserv Biol* **29**, 88-98, doi:10.1111/cobi.12370 (2015).

354 10 Turner, D. J., Smyth, A. K., Walker, C. M. & Lowe, A. J. in *Terrestrial Ecosystem Research Infrastructures*
355 341-368 (CRC Press, 2017).

356 11 Sullivan, B. L. *et al.* eBird: A citizen-based bird observation network in the biological sciences. *Biol*
357 *Conserv* **142**, 2282-2292, doi:10.1016/j.biocon.2009.05.006 (2009).

358 12 Rausher, M. D., McPeck, M. A., Moore, A. J., Rieseberg, L. & Whitlock, M. C. Data Archiving. *Evolution*
359 **64**, 603-604, doi:10.1111/j.1558-5646.2009.00940.x (2010).

360 13 Whitlock, M. C., McPeck, M. A., Rausher, M. D., Rieseberg, L. & Moore, A. J. Data Archiving. *Am Nat*
361 **175**, E45-146, doi:10.1086/650340 (2010).

362 14 Fuller, R. A., Lee, J. R. & Watson, J. E. M. Achieving open access to conservation science. *Conserv Biol*
363 **28**, 1550-1557 (2014).

364 15 Ngo, H. *et al.* First population assessment of two cryptic Tiger Geckos (*Goniurosaurus*) from northern
365 Vietnam: Implications for conservation. *Amphibian & Reptile Conservation* **10**, 34-45 (2016).

366 16 Lindenmayer, D. & Scheele, B. Do not publish. *Science* **356**, 800-801, doi:10.1126/science.aan1362
367 (2017).

368 17 Runge, C. A., Tulloch, A., Hammill, E., Possingham, H. P. & Fuller, R. A. Geographic range size and
369 extinction risk assessment in nomadic species. *Conserv Biol* **29**, 865-876 (2015).

370 18 Herzog, S. K., Maillard Z., O., Embert, D., Caballero, P. & Quiroga, D. Range size estimates of Bolivian
371 endemic bird species revisited: the importance of environmental data and national expert knowledge.
372 *Journal of Ornithology* **153**, 1189-1202, doi:10.1007/s10336-012-0850-2 (2012).

373 19 Chapman, A. D. & Grafton, O. Guide to best practices for generalising sensitive species occurrence
374 data. (Report for the Global Biodiversity Information Facility, Copenhagen, 2008).

- 375 20 Alonso, L. E., Deichmann, J. L., McKenna, S. A., Naskrecki, P. & Richards, S. J. (University of Chicago
376 Press, Chicago, 2011).
- 377 21 Reichman, O. J., Jones, M. B. & Schildhauer, M. P. Challenges and Opportunities of Open Data in
378 Ecology. *Science* **331**, 703-705, doi:10.1126/science.1197962 (2011).
- 379 22 Wallis, J. C., Rolando, E. & Borgman, C. L. If We Share Data, Will Anyone Use Them? Data Sharing and
380 Reuse in the Long Tail of Science and Technology. *Plos One* **8**, doi:10.1371/journal.pone.0067332
381 (2013).
- 382 23 Borgman, C. L. The Conundrum of Sharing Research Data. *J Am Soc Inf Sci Tec* **63**, 1059-1078,
383 doi:10.1002/asi.22634 (2012).
- 384 24 McKiernan, E. C. *et al.* The open research value proposition: How sharing data can help researchers
385 succeed. *Figshare*, doi.org/10.6084/m6089.figshare.1619902.v1619902, Retrieved 1619901 Nov
386 1612017, doi:doi.org/10.6084/m9.figshare.1619902.v2 (2015).
- 387 25 Borgman, C. L. *et al.* Knowledge infrastructures in science: data, diversity, and digital libraries. *Int J*
388 *Digit Librarie* **16**, 207-227, doi:10.1007/s00799-015-0157-z (2015).
- 389 26 Johnson, T. R. & van Densen, W. L. T. Benefits and organization of cooperative research for fisheries
390 management. *ICES Journal of Marine Science* **64**, 834-840, doi:10.1093/icesjms/fsm014 (2007).
- 391 27 Maxwell, S. L., Fuller, R. A., Brooks, T. M. & Watson, J. E. Biodiversity: The ravages of guns, nets and
392 bulldozers. *Nature* **536**, 143-145 (2016).
- 393 28 Fisher, D. O. Trajectories from extinction: where are missing mammals rediscovered? *Global Ecol*
394 *Biogeogr* **20**, 415-425, doi:10.1111/j.1466-8238.2010.00624.x (2011).
- 395 29 Fisher, D. O. & Blomberg, S. P. Correlates of rediscovery and the detectability of extinction in
396 mammals. *P Roy Soc B-Biol Sci* **278**, 1090-1097, doi:10.1098/rspb.2010.1579 (2011).
- 397 30 Kissling, W. D. *et al.* Building essential biodiversity variables (EBVs) of species distribution and
398 abundance at a global scale. *Biological Reviews* **93**, 600-625, doi:10.1111/brv.12359 (2018).
- 399 31 Proença, V. *et al.* Global biodiversity monitoring: From data sources to Essential Biodiversity Variables.
400 *Biol Conserv* **213**, 256-263, doi:10.1016/j.biocon.2016.07.014 (2017).

- 401 32 Trindade-Filho, J., de Carvalho, R. A., Brito, D. & Loyola, R. D. How does the inclusion of Data Deficient
402 species change conservation priorities for amphibians in the Atlantic Forest? *Biodivers Conserv* **21**,
403 2709-2718, doi:10.1007/s10531-012-0326-y (2012).
- 404 33 Setiawan, R. *et al.* Preventing Global Extinction of the Javan Rhino: Tsunami Risk and Future
405 Conservation Direction. *Conserv Lett* **11**, n/a-n/a, doi:10.1111/conl.12366 (2018).
- 406 34 Leader-Williams, N., Baldus, R. D. & Smith, R. J. in *Recreational hunting, conservation and rural*
407 *livelihoods: Science and practice* (eds B. Dickson, J. Hutton, & W.A. Adams) 296-316 (John Wiley &
408 Sons, 2009).
- 409 35 Loveridge, A. J., Hemson, G., Davidson, Z. & Macdonald, D. W. in *Wild felid biology and management*
410 (eds D.W. Macdonald & A.J. Loveridge) 283-304 (Oxford University Press, 2010).
- 411 36 Steven, R., Pickering, C. & Guy Castley, J. A review of the impacts of nature based recreation on birds. *J*
412 *Environ Manage* **92**, 2287-2294, doi:10.1016/j.jenvman.2011.05.005 (2011).
- 413 37 Gonzalez, J. A. Effects of Harvesting of Waterbirds and Their Eggs by Native People in the Northeastern
414 Peruvian Amazon. *Waterbirds: The International Journal of Waterbird Biology* **22**, 217-224,
415 doi:10.2307/1522210 (1999).
- 416 38 Arlettaz, R. *et al.* From Publications to Public Actions: When Conservation Biologists Bridge the Gap
417 between Research and Implementation. *Bioscience* **60**, 835-842, doi:10.1525/bio.2010.60.10.10
418 (2010).
- 419 39 Burgman, M. A. *Risks and decisions for conservation and environmental management*. (Cambridge
420 University Press, 2005).
- 421 40 Kelly, C. L., Pickering, C. M. & Buckley, R. C. Impacts of tourism on threatened plant taxa and
422 communities in Australia. *Ecological Management & Restoration* **4**, 37-44, doi:10.1046/j.1442-
423 8903.2003.00136.x (2003).
- 424 41 Ballantyne, M. & Pickering, C. Ecotourism as a threatening process for wild orchids. *Journal of*
425 *Ecotourism* **11**, 34-47, doi:10.1080/14724049.2011.628398 (2012).
- 426 42 Hissmann, K. *et al.* The South African coelacanths - an account of what is known after three
427 submersible expeditions : coelacanth research. *South African Journal of Science* **102**, 491-500 (2006).

428 43 Erdmann, M. Lessons learned from the conservation campaign for the Indonesian coelacanth,
429 Latimeria menadoensis : coelacanth research. *South African Journal of Science* **102**, 501-504 (2006).

430 44 Tulloch, V. J. D. *et al.* Why do we map threats? Linking threat mapping with actions to make better
431 conservation decisions. *Front Ecol Environ* **13**, 91-99, doi:10.1890/140022 (2015).

432 45 IUCN. The IUCN Red List of Threatened Species. Version 2017-3. <<http://www.iucnredlist.org>>.
433 (Downloaded on 05 January 2018, 2017).

434 46 Michener, W. K. Ecological data sharing. *Ecological Informatics* **29**, 33-44,
435 doi:<http://dx.doi.org/10.1016/j.ecoinf.2015.06.010> (2015).

436 47 Smith, M., Szongott, C., Henne, B. & Von Voigt, G. in *Digital Ecosystems Technologies (DEST), 2012 6th*
437 *IEEE International Conference on.* 1-6 (IEEE).

438 48 Mancini, A. & Koch, V. Sea turtle consumption and black market trade in Baja California Sur, Mexico.
439 *Endangered Species Research* **7**, 1-10 (2009).

440 49 Barry, C. Australia's wildlife blackmarket trade. *Australian Geographic* **August 16**,
441 doi:[http://www.australiangeographic.com.au/topics/wildlife/2011/08/australias-wildlife-blackmarket-](http://www.australiangeographic.com.au/topics/wildlife/2011/08/australias-wildlife-blackmarket-trade/)
442 [trade/](http://www.australiangeographic.com.au/topics/wildlife/2011/08/australias-wildlife-blackmarket-trade/) (2011).

443 50 Stiles, D., Redmond, I., Cress, D., Nellemann, C. & Formo, R. K. Stolen Apes: The Illicit Trade in
444 Chimpanzees, Gorillas, Bonobos and Orangutans. A Rapid Response Assessment. (United Nations
445 Environment Programme, GRID-Arendal. www.grida.no, 2013).

446 51 Auliya, M. *et al.* Trade in live reptiles, its impact on wild populations, and the role of the European
447 market. *Biol Conserv* **204, Part A**, 103-119, doi:<https://doi.org/10.1016/j.biocon.2016.05.017> (2016).

448 52 Collins-Kreiner, N., Malkinson, D., Labinger, Z. & Shtainvarz, R. Are birders good for birds? Bird
449 conservation through tourism management in the Hula Valley, Israel. *Tourism Management* **38**, 31-42,
450 doi:<http://dx.doi.org/10.1016/j.tourman.2013.01.009> (2013).

451 53 Lowe, A. J. *et al.* Publish openly but responsibly. *Science* **357**, 141-141, doi:10.1126/science.aao0054
452 (2017).

453 54 Michener, W. K., Brunt, J. W., Helly, J. J., Kirchner, T. B. & Stafford, S. G. Nongeospatial metadata for
454 the ecological sciences. *Ecol Appl* **7**, 330-342, doi:Doi 10.2307/2269427 (1997).

455 55 Piwowar, H. A., Day, R. S. & Fridsma, D. B. Sharing Detailed Research Data Is Associated with Increased
456 Citation Rate. *Plos One* **2**, e308, doi:ARTN e308 10.1371/journal.pone.0000308 (2007).

457 56 Campbell, E. G. & Bendavid, E. Data-sharing and data-withholding in genetics and the life sciences:
458 results of a national survey of technology transfer officers. *J Health Care Law Policy* **6**, 241-255 (2003).

459 57 Tenopir, C. *et al.* Data sharing by scientists: practices and perceptions. *Plos One* **6**, e21101,
460 doi:10.1371/journal.pone.0021101 (2011).

461 58 Keniger, L., Gaston, K., Irvine, K. & Fuller, R. What are the Benefits of Interacting with Nature?
462 *International Journal of Environmental Research and Public Health* **10**, 913 (2013).

463 59 Rosen, G. E. & Smith, K. F. Summarizing the Evidence on the International Trade in Illegal Wildlife.
464 *EcoHealth* **7**, 24-32, doi:10.1007/s10393-010-0317-y (2010).

465 60 Boakye, M. K., Pietersen, D. W., Kotzé, A., Dalton, D.-L. & Jansen, R. Knowledge and Uses of African
466 Pangolins as a Source of Traditional Medicine in Ghana. *Plos One* **10**, e0117199,
467 doi:10.1371/journal.pone.0117199 (2015).

468 61 Gong, S.-P., Chow, A. T., Fong, J. J. & Shi, H.-T. The chelonian trade in the largest pet market in China:
469 scale, scope and impact on turtle conservation. *Oryx* **43**, 213-216, doi:10.1017/S0030605308000902
470 (2009).

471 62 Stuart, B. L., Rhodin, A. G. J., Grismer, L. L. & Hansel, T. Scientific Description Can Imperil Species.
472 *Science* **312**, 1137-1137, doi:10.1126/science.312.5777.1137b (2006).

473 63 Purcell, S. W. *et al.* Sea cucumber fisheries: global analysis of stocks, management measures and
474 drivers of overfishing. *Fish and Fisheries* **14**, 34-59, doi:10.1111/j.1467-2979.2011.00443.x (2013).

475 64 Packer, C. *et al.* Effects of Trophy Hunting on Lion and Leopard Populations in Tanzania. *Conserv Biol*
476 **25**, 142-153, doi:10.1111/j.1523-1739.2010.01576.x (2011).

477 65 Williams, V. L., Cunningham, A. B., Kemp, A. C. & Bruyns, R. K. Risks to Birds Traded for African
478 Traditional Medicine: A Quantitative Assessment. *Plos One* **9**, e105397,
479 doi:10.1371/journal.pone.0105397 (2014).

480 66 Meeuwig, J. J., Harcourt, R. G. & Whoriskey, F. G. When Science Places Threatened Species at Risk.
481 *Conserv Lett* **8**, 151-152, doi:10.1111/conl.12185 (2015).

482 67 Crawford, A. & Bernstein, J. MEAs, Conservation and Conflicts. A case study of Virunga National Park,
483 DRC. (International Institute for Sustainable Development, Geneva, Switzerland, 2008).

484 68 Jenkins, M. Who murdered the Virunga gorillas. *National Geographic* **214**, 34-65 (2008).

485 69 Sattler, P. & Williams, R. The Conservation Status of Queensland's Bioregional Ecosystems. (QLD
486 Environment Protection Agency, Brisbane, Australia, 1999).

487 70 Kearney, M. R., Porter, W. P. & Murphy, S. A. An estimate of the water budget for the endangered
488 night parrot of Australia under recent and future climates. *Climate Change Responses* **3**, 14,
489 doi:10.1186/s40665-016-0027-y (2016).

490 71 Murphy, S. A., Silcock, J., Murphy, R., Reid, J. & Austin, J. J. Movements and habitat use of the night
491 parrot *Pezoporus occidentalis* in south-western Queensland. *Austral Ecology* **42**, 858-868,
492 doi:10.1111/aec.12508 (2017).

493 72 Sadovy de Mitcheson, Y. *et al.* Fishing groupers towards extinction: a global assessment of threats and
494 extinction risks in a billion dollar fishery. *Fish and Fisheries* **14**, 119-136, doi:10.1111/j.1467-
495 2979.2011.00455.x (2013).

496 73 Jumin, R. *et al.* From Marxan to management: ocean zoning with stakeholders for Tun Mustapha Park
497 in Sabah, Malaysia. *Oryx*, 1-12, doi:10.1017/S0030605316001514 (2017).

498 74 Adams, V. M., Tulloch, V. J. & Possingham., H. P. Land-sea conservation assessment for Papua New
499 Guinea: A report on the work undertaken to fulfil the terms of the project Review and Integration of
500 the Terrestrial and Marine Program of Works on Protected Areas. (University of Queensland, Australia,
501 and Conservation and Environment Protection Authority, Papua New Guinea, 2016).

502 75 Government of Papua New Guinea. National Marine Conservation Assessment for Papua New Guinea.
503 (Conservation and Environment Protection Authority, Papua New Guinea, 2015).

504 76 Hamilton, R., de Mitcheson, Y. S. & Aguilar-Perera, A. The Role of Local Ecological Knowledge in the
505 Conservation and Management of Reef Fish Spawning Aggregations. *Fish Fish Ser* **35**, 331-369,
506 doi:10.1007/978-94-007-1980-4_10 (2012).

507 77 Lavery, T. H. & Judge, H. A new species of giant rat (Muridae, *Uromys*) from Vangunu, Solomon Islands.
508 *Journal of Mammalogy*, gyx116-gyx116, doi:10.1093/jmammal/gyx116 (2017).

- 509 78 Barcelona, J. F., Pelsner, P. B., Balette, D. S. & Co, L. L. Taxonomy, ecology, and conservation status of
510 Philippine Rafflesia (Rafflesiaceae). *Blumea - Biodiversity, Evolution and Biogeography of Plants* **54**, 77-
511 93, doi:10.3767/000651909X474122 (2009).
- 512 79 Olson, R. J. & McCord, R. A. in *Ecological Data: Design, Management, and Processing* (ed J.W. Brunt
513 W.K. Michener) 117-141 (Blackwell Science Ltd., 2000).

514

515 **Acknowledgements**

516 AITT was supported by an Australian Research Council Discovery Early Career Researcher Award DE170100599.
517 EB, GE, NPL and LR were supported by the Australian Government National Environmental Science
518 Programme's Threatened Species Recovery Hub. NPL was partially funded by Bush Heritage Australia. Richard
519 Alcorn (eBird), Tanya Laity (Australian Government Department of the Environment and Energy), Steve
520 Murphy and Alex Kutt (Bush Heritage Australia) provided feedback on early drafts. Jessica Miller and Richard
521 Fuller contributed to early discussions.

522

523 **Author contributions:** AITT led the development of the risk assessment and decision tree with contributions
524 from all authors to the final decision protocol. All authors provided ideas and critical feedback and co-wrote
525 the manuscript.

526

527 **Competing interests:** The authors have no competing financial or non-financial interests.

528

529 **Data availability:** Data sharing not applicable to this article as no datasets were generated or analysed during
530 the current study.

531

532 **Table 1.** Likely objectives of various data owners (potential data providers) for sharing biodiversity data and
 533 perceptions of how publishing species data might advance or undermine that objective.

Objective for sharing data	Examples of data authors	Perception of how publishing data advances objective	Perception of how publishing data undermines objective
<i>Minimise risk of species extinction</i>	Government agencies, NGO ecologists, scientists, land managers	Avoid impacts and increase protection in known locations; Increase species ecological knowledge; Inform status assessments (e.g. IUCN Red List).	Increase human access to species' locations, leading to poaching ⁵¹ , disease spread, habitat degradation (trampling, alien species spread) or disturbance ^{36,52} .
<i>Maximise ability to get funding for biodiversity management</i>	NGO ecologists, scientists, land managers	Secure future funding by demonstrating importance and effectiveness of project (e.g. important species detected).	Reduce future funding by disproving importance or effectiveness of project (e.g. important species not detected), or exposing inappropriately collected data; Hinder projects and profitability by inspiring extra regulation.
<i>Inform and improve future monitoring</i>	Government agencies at local, state or national levels	Evaluate progress towards conservation objectives (e.g. identify trends); Develop spatial units for data aggregation and future sampling.	Violate privacy of stakeholders granting access to study area (e.g. landholders, indigenous groups), limiting future access.
<i>Maximise data reliability for improving knowledge</i>	IUCN, government agencies, scientists	Decrease false negative error rate; Illuminate gaps in knowledge of species; Reduce data deficiency for species ⁵³ .	Increase false positive error rate if datasets vary in rigour or accuracy, wasting effort on incorrect records.
<i>Minimise data loss/ entropy and survey redundancy</i>	Museums, developers, consultants, government agencies	Reduce risk of losing access to data and legacy effects of loss of local champion ⁵⁴ ; Maximise efficiency of limited resources to collect structured data.	Increase visitation to private study or recreational locations, reducing novelty or utility of data; Reduce knowledge power and competitive edge (e.g. to win survey tenders through local expertise).
<i>Maximise academic output or impact</i>	Scientific researcher in academic institution	Increase collaborations, citations and use of research ⁵⁵ ; Verify published results.	Others use data without acknowledging/ citing source ^{56,57} ; Opportunity cost (e.g. time) of compiling/sharing data; Embarrassment if others find errors in analysis (e.g. publish rebuttals).
<i>Maximise public interest in biodiversity</i>	Citizen science programs, citizen scientists, government agencies	Engage the public in nature; Human health benefits of wildlife access ⁵⁸ .	Increase human access to species' locations, leading to inadvertent habitat degradation or disturbance and reduced site value ^{36,52} .

534 **Table 2. Threats to species related to sharing of biodiversity data.** Making localities of some species available
535 to the public can increase one or more of three major threats to species: poaching (driven by illegal wildlife
536 trade), hunting/collecting for food, medicine and trophies, and habitat disturbance and degradation. Effects on
537 populations of sensitive species range from disturbance of regular behaviours and reduced reproductive rates
538 to reduced population survival and possible local or global extinctions, but there is little direct evidence linking
539 data publication to these impacts. Literature review conducted by searching Web of Science (20/8/2017) for
540 keywords “species” AND “data publication” or “data sharing” or “publication of data” AND “threat” or
541 “poach*” or “wildlife trade” or “recreational hunting” or “trophy hunting”.

Threat	Examples of the scale of the threat	Direct evidence of data publication leading to population impacts
<i>Collection for international wildlife markets</i>	Mortality and associated population declines due to illegal trafficking of ivory (~5370 dead elephants) between 1996-2008 ⁵⁹ .	Published data (digital archives, scientific journals, social media) led to poachers collecting 1000s of South African succulents to sell to European plant enthusiasts ^a .
<i>Collection for local traditional medicine</i>	Population declines of African pangolins due to harvesting scales/bones for spiritual healing ⁶⁰ .	No published literature found directly linking data publication to impacts.
<i>Collection for pet trade</i>	Pet market trade in CITES-listed turtle species in China ⁶¹ ; illegal import of >20 million live reptiles to EU between 2004-2014 ⁵¹ .	Local extinctions of newly discovered restricted-range reptiles of value in pet markets following formal publications of locality information ⁶² .
<i>Resource use such as harvesting animals or eggs for food</i>	Declines in marine stocks (e.g. sea cucumbers, tuna) due to over-fishing ⁶³ ; reduced nesting and colony abandonment of Andes flamingos and Peruvian heronries due to egg harvest ³⁷ .	No published literature found directly linking data publication to impacts. Anecdotal records of resource declines (e.g. in Coral Triangle spawning locations).
<i>Recreational hunting</i>	Areas with highest trophy hunting levels have highest population declines of Tanzanian lions and leopards ⁶⁴ .	Reduced population survival and group cohesion of lions in Africa due to corrupt tourist hunting practices tracking known individuals/prides ^{34,35} .
<i>Cultural beliefs about negative impacts of species</i>	Illegal killing and disturbance of native flying foxes in Australia perceived as a nuisance due to noise, smell and droppings at roost sites ^d .	Research data tracking endangered white sharks used by Western Australian government agency to kill sharks in attempt to reduce shark-human interactions ⁶⁶ .
<i>Recreational disturbance for wildlife watching</i>	Habitat degradation and disturbance of animal behaviours by bird watchers in popular bird-watching locations ³⁶ .	Data shared on breeding location of painted snipe pair resulted in nest being abandoned due to disturbance by wildlife enthusiasts ^b .
<i>Killing/ habitat destruction to</i>	Killing of mountain gorillas in DRC allegedly by rebels, soldiers, corrupt officials to discredit	'Panic clearing' by Australian landholders of vegetation needing protective mechanisms

prevent conservationists and facilitate access to World after government report⁶⁹ mapped
conservation Heritage region for illegal charcoal trade^{67,68} occurrence of rare/threatened communities^c.

- 542 a. <http://e360.yale.edu/features/unnatural-surveillance-how-online-data-is-putting-species-at-risk>
- 543 b. <http://www.shanghaibirding.com/2017/08/24/painted-snipe-ethics>
- 544 c. <https://theconversation.com/australia-is-a-global-top-ten-deforester-and-queensland-is-leading-the-way-87259>
- 545 d. <https://www.animalecologylab.org/not-in-my-backyard.html>

546

547 **Table 3.** Examples of how decision tree can be used to support decisions to share data (see Supplementary
 548 Table 2 for details).

Species and location	Risks of data sharing	Conservation mechanisms and benefits of data publication	Decision tree action and reasons
1. Night Parrot <i>Pezoporus occidentalis</i> , arid zone of Australia	No population of live birds known until 2013. Exact location withheld from the public. To limit risks of poaching or disturbance from birdwatchers very few conservation professionals, government representatives and researchers had access to location data and calls.	In 2016, Queensland State Government made discovery location an exclusion zone with imprisonment if accessed unlawfully. Bird’s call and data on habitat preferences released when conservation mechanisms were in place. At least three new populations discovered since by people using recorded call ^{70,71} .	Species was at risk of exploitation due to <i>in situ</i> activities (disturbance due to human activities) in 2013. Decision: MASK SPECIES LOCATIONS, NOT IDENTITIES In 2018 protection mechanisms are in place to mitigate declines. Decision: PUBLIC DATA
2. Fish spawning aggregations, Pacific Ocean	Species that spawn seasonally in large groups are easy for fishers to locate and exploit. Over-exploitation of reef fish spawning aggregations led to collapse of several species following discovery of new aggregation locations, with 20 species of grouper at risk of extinction if current overfishing trends continue.	Controlled release of spawning data to researchers by the NGO SCRFA (Science and Conservation of Fish Aggregations) through data sharing confidentiality agreements enabled decision-makers to target spatial conservation planning to protect species such as threatened grouper (Serranidae), snapper (Lutjanidae) and emperor fish (Lethrinidae) ⁷²⁻⁷⁶ .	Species is at risk of <i>ex situ</i> exploitation for food and fish trade. Conservation/policy mechanisms to mitigate declines not in place at all sites. Location data useful to inform management. Data can be requested by researchers and used under protocols that restrict publishing of maps identifying spawning locations. Decision: RESTRICT DATA: MASK SPECIES IDENTITIES AND LOCATIONS

<p>3. Vangunu Giant Rat <i>Uromys vika</i>, Solomon Islands</p>	<p>Recently discovered mammal known from a single specimen on Vangunu Island⁷⁷. Location is remote and difficult to access, requiring permission from landowners and tribal chiefs. Forest inhabited by <i>U. vika</i> is threatened by logging and not secured under any formal conservation land.</p>	<p>Local community attempting to protect habitat as the Zaira Community Resource Management Area (ZCRMA), which also supports a vulnerable bat <i>Pteralopex taki</i> and nesting leatherback turtles <i>Dermochelys coriacea</i>. It is hoped that increased recognition will attract further support for ZCRMA and thus protect <i>U. vika</i>.</p>	<p>Species has no <i>ex situ</i> economic value and not threatened by disturbance from increased visitation due to remoteness of location. Publication of holotype location data will increase recognition of forests' conservation value and may help conservation efforts to set up protected areas.</p> <p>Decision: PUBLIC DATA</p>
<p>4. Parasitic corpse flower <i>Rafflesia</i>, Malaysia</p>	<p>Populations consist of few individuals, are relatively accessible and threatened by logging, agriculture and disturbance by unsustainable ecotourism. Releasing species occurrence data without conservation measures would increase risk of decline of <i>Rafflesia</i> and host plants through increased visitation⁷⁸.</p>	<p>Visitors to <i>Rafflesia</i> blooming events provide nature ecotourism revenue. Populations occur mostly outside protected areas and there are no proposed protection mechanisms or policies to mitigate declines. Data might be used by conservation planners to propose new protected areas.</p>	<p>Individuals have <i>in situ</i> value for tourism, but could be threatened by increased visitation. Publication of location data is needed to identify sites with conservation value, but human traffic must be managed.</p> <p>Decision: RESTRICT DATA: MASK SPECIES IDENTITIES NOT LOCATIONS.</p>

Box 1. Glossary for decision tree

Access: whether individual/population can be physically reached by humans, restrictive access can be due either due to natural processes/factors (e.g., geological = species located on a sea cliff), or human intervention (e.g., protection = species in a strictly controlled site).

Associated species: non-target species that may be dependent on the target species, for example, due to physical/structural relationships (e.g., liana on a tropical tree), or nutritional interactions (e.g., obligate nectarivore or pollen-producer, or obligate seed disperser such as flying fox).

Conservation/policy mechanisms: actions that aim to improve species/population persistence by mitigating threats causing population declines.

Data repository or archive: a permanent collection of data sets with accompanying metadata usually stored in an online cloud service so that a variety of users can readily access, understand, download and use data ⁷⁹.

Ex situ economic value: individuals or populations have monetary worth outside of natural/native site, e.g., exploitation through harvesting of individuals for food, medicine or wildlife trade.

Impact on other species: for example, ground-level species may be (inadvertently) trampled/damaged by people walking to access the target species, or arboreal species by people climbing trees to view another target species.

Impact on population: direct (removal), physical (damage to individuals), or environmental (habitat degradation/reduction) impact, such that population viability is compromised.

In situ value: individuals/populations have monetary or cultural worth in natural/native site, e.g., for ecotourism.

In situ disturbance: impacts on individuals/populations in natural/native site due to human activity, e.g., invasive species, tourism, habitat degradation.

Public data: information on species locations, abundance, ecology published with no restrictions so that it is openly available to the public (i.e. primary data with no denaturing or masking).

Restrict data: information on species locations, abundance, ecology only published after being masked, generalised or converted to habitat suitability model. Some masked species and location data are likely to be stored offline but models may be publicly available.